arXiv:1304.0858v1 [nlin.SI] 3 Apr 2013

Hypergeometric solutions to the symmetric q-Painlevé equations

Kenji Kajiwara[1] and Nobutaka Nakazono[2]

[1]: Institute of Mathematics for Industry, Kyushu University,
744 Motooka, Fukuoka 819-8581, Japan.
kaji@imi.kyushu-u.ac.jp
[2]: Graduate School of Mathematics, Kyushu University,
744 Motooka, Fukuoka 819-8581, Japan.
n-nakazono@kyudai.jp

April 4, 2013

Abstract

We consider the symmetric q-Painlevé equations derived from the birational representation of affine Weyl groups by applying the projective reduction and construct the hypergeometric solutions. Moreover, we discuss continuous limits of the symmetric q-Painlevé equations to Painlevé equations together with their hypergeometric solutions.

2010 Mathematics Subject Classification: 33D15, 33E17, 34M55, 39A13
Keywords and Phrases: q-Painlevé equation; Painlevé equation; basic hypergeometric function; hypergeometric function; saddle point method; projective reduction; continuous limit

1 Introduction

The discrete Painlevé equations, together with the Painlevé equations, are now widely recognized as the one of the most important families of the integrable systems (see, for example, [7]). Originally, the discrete Painlevé equations were discovered in the form of single second-order difference equations and identified as the discrete analogue of the Painlevé equation [1–3, 33, 38] Then they were generalized to simultaneous first-order equations by a singularity confinement criterion [8, 38]. A typical example is the following equation known as a discrete Painlevé II equation [33, 38]:

$$x_{n+1} + x_{n-1} = \frac{(an+b)x_n + c}{1 - {x_n}^2}, \tag{1.1}$$

where x_n is the dependent variable, n is the independent variable, and a, b, $c \in \mathbb{C}$ are parameters. We note that (1.1) was found as the equation satisfied by the constant terms of the orthogonal polynomials on the unit circle [33]. By applying the singularity confinement criterion, (1.1) is generalized to

$$x_{n+1} + x_{n-1} = \frac{(an+b)x_n + c + (-1)^n d}{1 - {x_n}^2}, \tag{1.2}$$

where d is a parameter, with its integrability preserved. Introducing the dependent variables X_n and Y_n by

$$X_n = x_{2n}, \quad Y_n = x_{2n-1}, \tag{1.3}$$

then (1.2) can be rewritten as

$$Y_{n+1} + Y_n = \frac{(2an+b)X_n + c + d}{1 - {X_n}^2}, \quad X_{n+1} + X_n = \frac{(a(2n+1)+b)Y_{n+1} + c - d}{1 - {Y_{n+1}}^2}. \tag{1.4}$$

Equation (1.4) is known as a discrete Painlevé III equation since it admits a continuous limit to the Painlevé III equation [4, 30, 40]. Conversely, (1.1) can be recovered from (1.4) by putting $d = 0$ and (1.3). This procedure is referred to as "*symmetrization*" of (1.4), which comes from the terminology of the Quispel–Roberts–Thompson (QRT) mapping [35, 36]. After this terminology, (1.4) is sometimes called the "*asymmetric*" discrete Painlevé II equation, and (1.1) is called the "*symmetric*" discrete Painlevé III equation [21].

It appears as though the symmetrization is a simple specialization on the level of the equation, but the following problems were known: (i) According to Sakai's theory [42], the Painlevé and discrete Painlevé equations are classified by the underlying space of initial conditions. Moreover, the discrete Painlevé equations arise as the birational mappings corresponding to the translations of the affine Weyl groups associated with the space of initial conditions. The asymmetric discrete Painlevé equations are characterized in this manner, however, it was not known how to characterize the symmetric equations as the action of affine Weyl groups. (ii) The Painlevé and discrete Painlevé equations admit the particular solutions expressible in terms of the hypergometric type functions (hypergeometric solutions) when some of the parameters take special values (see, for example, [14, 15] and references therein). However, the hypergeometric solutions to the symmetric discrete Painlevé equation cannot be obtained by the naïve specialization of those to the corresponding asymmetric equation. For example, (1.1) has the hypergeometric solution expressible in terms of the parabolic cylinder function (Weber function) [13, 18]. On the other hand, (1.4) admits the hypergeometric solution in terms of the confluent hypergeometric function [16, 22, 32]. The crucial point is that although the former function is expressed as a specialization of the latter, this specialization is not consistent with the symmetrization.

In [16], the mechanism of the symmetrization was investigated in detail by taking an example of q-Painlevé equation with the affine Weyl group symmetry of type $(A_2 + A_1)^{(1)}$. Then it was shown that in general, various discrete dynamical systems of Painlevé type can be obtained from elements of infinite order that are not necessarily translations in the affine Weyl group by taking the projection on appropriate subspaces of the parameter spaces. Such a procedure is called a *projective reduction*, and the symmetrization can be understood as a kind of projective reduction. Moreover, the above nontrivial inconsistency among the hypergeometric solutions are explained by the factorization of the linear difference operators associated with the three-term relation of the hypergeometric functions.

In spite of understanding the mechanism of the symmetrization or the projective reduction, it is still nontrivial to construct the hypergeometric solutions to the symmetric discrete Painlevé equations. In particular, since the continuous limit of the symmetric discrete Painlevé equation is different from the corresponding asymmetric one, it is important to construct the hypergeometric solutions to symmetric equations and consider their continuous limits.

The purpose of this paper is to construct the simplest hypergeometric solutions to the symmetric q-Painlevé equations. In [14, 15] the simplest hypergeometric solutions to all possible q-Painlevé equations in the Sakai's list have been constructed. Similarly, in this paper, we present the list of the simplest hypergeometric solutions to the symmetric q-Painlevé equations reduced from the asymmetric ones with the affine Weyl group symmetry of type $E_8^{(1)}$, $E_7^{(1)}$, $E_6^{(1)}$, $D_5^{(1)}$, $A_4^{(1)}$ and $(A_2 +$

$A_1)^{(1)}$. We also aim to consider the continuous limits of those symmetric q-Painlevé equations to the Painlevé equations, together with their hypergeometric solutions.

This paper is organized as follows: in Section 2, we show the symmetric q-Painlevé equations by applying the projective reduction to q-P($(A_2+A_1)^{(1)}$), q-P($A_4^{(1)}$), q-P($D_5^{(1)}$), q-P($E_6^{(1)}$), q-P($E_7^{(1)}$) and q-P($E_8^{(1)}$), where the q-Painlevé equation with the affine Weyl group of type X is denoted by q-P(X). In Section 3, we construct the hypergeometric solutions to the symmetric q-Painlevé equations derived in Section 2. In Section 4, we discuss the continuous limits of the symmetric q-Painlevé equations and their hypergeometric solutions. In Section 5, we prove that hypergeometric function appearing in the hypergeometric solution to the symmetric q-P($A_4^{(1)}$) actually reduces to the Weber function by applying the saddle point method to its integral representation. Some concluding remarks are given in Section 6.

2 Symmetric q-Painlevé equations

In this section, we apply the projective reduction to the q-Painlevé equations q-P($(A_2+A_1)^{(1)}$), q-P($A_4^{(1)}$), q-P($D_5^{(1)}$), q-P($E_6^{(1)}$), q-P($E_7^{(1)}$) and q-P($E_8^{(1)}$), respectively, to obtain their symmetric forms. In the following, we use the notations

$$\begin{aligned} &\overline{t}=qt, \quad \underline{t}=q^{-1}t, \quad f=f(t), \quad \overline{f}=f(\overline{t}), \quad \underline{f}=f(\underline{t}),\\ &p^2=q, \quad \tilde{t}=pt, \quad \utilde{t}=p^{-1}t, \quad \widetilde{f}=f(\tilde{t}), \quad \utilde{f}=f(\utilde{t}). \end{aligned} \tag{2.1}$$

Type $(A_2+A_1)^{(1)}$

q-P($(A_2+A_1)^{(1)}$) is given by [14, 15, 21, 39]

$$\overline{g}g=qc^2\frac{1+tf}{(t+f)f}, \quad \overline{f}f=qc^2\frac{1+a_2t\overline{g}}{(a_2t+\overline{g})\overline{g}}. \tag{2.2}$$

The continuous limit yields the Painlevé III equation. Applying the projective reduction with

$$f(t)=X(t), \quad g(t)=X(p^{-1}t), \quad a_2=p, \tag{2.3}$$

we obtain the symmetric q-P($(A_2+A_1)^{(1)}$) [16, 25, 27, 37]

$$\widetilde{X}\utilde{X}=p^2c^2\frac{1+tX}{(t+X)X}. \tag{2.4}$$

The continuous limit yields the Painlevé II equation.

Type $A_4^{(1)}$

q-P($A_4^{(1)}$) [43]:

$$(\overline{g}f-1)(gf-1)=t^2\frac{(f+a_1)(f+a_1{}^{-1})}{f+a_2t}, \quad (gf-1)(g\underline{f}-1)=q^{-1}t^2\frac{(g+a_1)(g+a_1{}^{-1})}{g+a_3t}. \tag{2.5}$$

Symmetric q-P($A_4^{(1)}$) [37, 43]:

$$(\widetilde{X}X - 1)(X\underset{\sim}{X} - 1) = t^2 \frac{(X + a_1)(X + a_1{}^{-1})}{X + a_2 t}\,, \tag{2.6}$$

where

$$f(t) = X(t), \quad g(t) = X(p^{-1}t), \quad a_3 = p^{-1}a_2. \tag{2.7}$$

The continuous limits of (2.5) and (2.6) yield the Painlevé V and IV equations, respectively.

Type $D_5^{(1)}$

q-P($D_5^{(1)}$) [11, 14, 39, 41]:

$$\overline{g}g = \frac{(f - q^{1/2}a_1 t)(f - q^{1/2}a_1{}^{-1}t)}{(f - a_2)(f - a_2{}^{-1})}, \qquad f\underline{f} = \frac{(g - a_3 t)(g - a_3{}^{-1}t)}{(g - a_4)(g - a_4{}^{-1})}. \tag{2.8}$$

Symmetric q-P($D_5^{(1)}$) [38]:

$$\widetilde{X}\underset{\sim}{X} = \frac{(X - a_1\tilde{t})(X - a_1{}^{-1}\tilde{t})}{(X - a_2)(X - a_2{}^{-1})}, \tag{2.9}$$

where

$$f(t) = X(t), \quad g(t) = X(p^{-1}t), \quad a_3 = a_1, \quad a_4 = a_2. \tag{2.10}$$

The continuous limits of (2.8) and (2.9) yield the Painlevé VI and III equations, respectively.

Type $E_6^{(1)}$

q-P($E_6^{(1)}$) [14, 15, 24, 39]:

$$\begin{cases} (\overline{g}f - 1)(gf - 1) = t^2 \dfrac{(f - b_1)(f - b_2)(f - b_3)(f - b_4)}{(f - b_5 t)(f - b_5{}^{-1}t)}, \\ (gf - 1)(g\underline{f} - 1) = q^{-1}t^2 \dfrac{(g - b_1{}^{-1})(g - b_2{}^{-1})(g - b_3{}^{-1})(g - b_4{}^{-1})}{(g - b_6 u)(g - b_6{}^{-1}u)}, \end{cases} \tag{2.11}$$

where

$$b_1 b_2 b_3 b_4 = 1, \quad u = p^{-1}t. \tag{2.12}$$

Symmetric q-P($E_6^{(1)}$) [38]:

$$(\widetilde{X}X - 1)(X\underset{\sim}{X} - 1) = t^2 \frac{(X - b_1)(X - b_1{}^{-1})(X - b_3)(X - b_3{}^{-1})}{(X - b_5 t)(X - b_5{}^{-1}t)}. \tag{2.13}$$

where

$$f(t) = X(t), \quad g(t) = X(p^{-1}t), \quad b_1 b_2 = 1, \quad b_3 b_4 = 1, \quad b_5 b_6 = 1. \tag{2.14}$$

Direct continuous limit of (2.11) to the Painlevé equations does not exist, while that of (2.13) yields the Painlevé V equation.

Type $E_7^{(1)}$

q-P($E_7^{(1)}$) [14, 15, 39]:

$$\begin{cases} \dfrac{(\overline{g}f-\overline{t}t)(gf-t^2)}{(\overline{g}f-1)(gf-1)} = \dfrac{(f-b_1t)(f-b_2t)(f-b_3t)(f-b_4t)}{(f-b_5)(f-b_6)(f-b_7)(f-b_8)}, \\ \dfrac{(gf-t^2)(g\underline{f}-t\underline{t})}{(gf-1)(g\underline{f}-1)} = \dfrac{(g-b_1{}^{-1}t)(g-b_2{}^{-1}t)(g-b_3{}^{-1}t)(g-b_4{}^{-1}t)}{(g-b_5{}^{-1})(g-b_6{}^{-1})(g-b_7{}^{-1})(g-b_8{}^{-1})}, \end{cases} \tag{2.15}$$

where

$$b_1b_2b_3b_4 = q, \quad b_5b_6b_7b_8 = 1. \tag{2.16}$$

Symmetric q-P($E_7^{(1)}$) [6]:

$$\frac{(\widetilde{X}X-\tilde{t}^2)(X\utilde{X}-t^2)}{(\widetilde{X}X-1)(X\utilde{X}-1)} = \frac{(X-b_1t)(X-b_1{}^{-1}\tilde{t})(X-b_3t)(X-b_3{}^{-1}\tilde{t})}{(X-b_5)(X-b_5{}^{-1})(X-b_7)(X-b_7{}^{-1})}. \tag{2.17}$$

where

$$f(t) = X(t), \quad g(t) = X(p^{-1}t), \quad b_1b_2 = p, \quad b_3b_4 = p, \quad b_5b_6 = 1, \quad b_7b_8 = 1. \tag{2.18}$$

Direct continuous limit of (2.15) to the Painlevé equations does not exist, while that of (2.17) yields the Painlevé VI equation.

Type $E_8^{(1)}$

q-P($E_8^{(1)}$) [14, 15, 24, 29, 39]:

$$\begin{cases} \dfrac{(\overline{g}\,\overline{u}t-f)(gut-f)-(\overline{u}^2t^2-1)(u^2t^2-1)}{(\overline{u}^{-1}t^{-1}\overline{g}-f)(u^{-1}t^{-1}g-f)-(1-\overline{u}^{-2}t^{-2})(1-u^{-2}t^{-2})} = \dfrac{P(f,t,m_1,\cdots,m_7)}{P(f,t^{-1},m_7,\cdots,m_1)}, \\ \dfrac{(fut-g)(\underline{f}u\underline{t}-g)-(u^2t^2-1)(u^2\underline{t}^2-1)}{(u^{-1}t^{-1}f-g)(u^{-1}\underline{t}^{-1}\underline{f}-g)-(1-u^{-2}t^{-2})(1-u^{-2}\underline{t}^{-2})} = \dfrac{P(g,u,m_7,\cdots,m_1)}{P(g,u^{-1},m_1,\cdots,m_7)}, \end{cases} \tag{2.19}$$

where

$$\begin{aligned} P(f,t,m_1,\cdots,m_7) = {} & f^4-m_1tf^3+(m_2t^2-3-t^8)f^2 \\ & +(m_7t^7-m_3t^3+2m_1t)f+t^8-m_6t^6+m_4t^4-m_2t^2+1. \end{aligned} \tag{2.20}$$

Here m_j ($j = 1,\cdots,8$) are the elementary symmetric functions of j-th degree in b_k ($k = 1,\cdots,8$), $m_8 = 1$, and $u = p^{-1}t$. Symmetric q-P($E_8^{(1)}$):

$$\frac{(\tilde{t}t\widetilde{X}-X)(t\utilde{t}\utilde{X}-X)-(\tilde{t}^2t^2-1)(t^2\utilde{t}^2-1)}{(\tilde{t}^{-1}t^{-1}\widetilde{X}-X)(t^{-1}\utilde{t}^{-1}\utilde{X}-X)-(\tilde{t}^{-2}t^{-2}-1)(t^{-2}\utilde{t}^{-2}-1)} = \frac{P(X,t,m_1,m_2,m_3,m_4,m_3,m_2,m_1)}{P(X,t^{-1},m_1,m_2,m_3,m_4,m_3,m_2,m_1)}. \tag{2.21}$$

where

$$f(t) = X(t), \quad g(t) = X(p^{-1}t), \quad m_i = m_{8-i} \quad (i = 1,2,3). \tag{2.22}$$

Direct continuous limits of (2.19) and (2.21) to the Painlevé equations are not known.

3 Hypergeometric solutions to the symmetric q-Painlevé equations

In this section, we construct the hypergeometric solutions to the symmetric q-Painlevé equations. We use the following conventions of q-analysis [5]. The basic hypergeometric series ${}_s\varphi_r$ is defined by

$$ {}_s\varphi_r\left(\begin{matrix}a_1,\cdots,a_s\\ b_1,\cdots,b_r\end{matrix};q,z\right)=\sum_{n=0}^{\infty}\frac{(a_1,\cdots,a_s;q)_n}{(b_1,\cdots,b_r;q)_n(q;q)_n}\left[(-1)^n q^{n(n-1)/2}\right]^{1+r-s} z^n, \tag{3.1}$$

where

$$(a_1,\cdots,a_s;q)_n=\prod_{i=1}^{s}(a_i;q)_n,\quad (a;q)_k=\prod_{i=1}^{k}(1-aq^{i-1}), \tag{3.2}$$

are the q-shifted factorials. The following special case of the basic hypergeometric series is known as the very-well-poised basic hypergeometric series

$$ {}_{r+1}W_r(a_1;a_4,a_5,\cdots,a_{r+1};q,z)={}_{r+1}\varphi_r\left(\begin{matrix}a_1,qa_1{}^{1/2},-qa_1{}^{1/2},a_4,\cdots,a_{r+1}\\ a_1{}^{1/2},-a_1{}^{1/2},qa_1a_4{}^{-1},\cdots,qa_1a_{r+1}{}^{-1}\end{matrix};q,z\right). \tag{3.3}$$

We also use the Jacobi theta function

$$\Theta(a;q)=(a;q)_\infty(qa^{-1};q)_\infty, \tag{3.4}$$

which satisfies the q-difference equation

$$\Theta(qa;q)=-a^{-1}\Theta(a;q). \tag{3.5}$$

3.1 Symmetric q-P$((A_2+A_1)^{(1)})$

Proposition 3.1 *Symmetric q-P$((A_2+A_1)^{(1)})$ (2.4) admits the hypergeometric solution*

$$\widetilde{X}=p^{1/2}\frac{\widetilde{G}}{G}, \tag{3.6}$$

$$G=Ae^{(\pi i/2)(\log t/\log p)}{}_1\varphi_1\left(\begin{matrix}0\\ -p\end{matrix};p,ip^{3/2}t\right)+Be^{(-\pi i/2)(\log t/\log p)}{}_1\varphi_1\left(\begin{matrix}0\\ -p\end{matrix};p,-ip^{3/2}t\right), \tag{3.7}$$

with

$$c=1. \tag{3.8}$$

Here, A and B are quasi-constants satisfying $A(t)=A(pt)$ and $B(t)=B(pt)$, respectively.

Proof. It is easily verified that (2.4) admits a specialization to the discrete Riccati equation

$$\widetilde{X}=-p\frac{1+tX}{X}, \tag{3.9}$$

when $c=1$. Then putting as (3.6), (3.9) is linearized to the three-term relation for G

$$\widetilde{\widetilde{G}}+p^{3/2}t\,\widetilde{G}+G=0. \tag{3.10}$$

Moreover, substituting the power series expression

$$G=\sum_{n=0}^{\infty}C_n t^{n+\rho},\qquad (\rho,\ C_n\in\mathbb{C}), \tag{3.11}$$

into (3.10), we obtain (3.7). □

In the following, we present the hypergeometric solutions to other cases. Since the results are verified by direct calculations, we omit the proof showing only the discrete Riccati equations and the linearized three-term relations. We also assume that A and B are quasi-constants.

3.2 Symmetric q-P($A_4^{(1)}$)

Proposition 3.2 *Symmetric* q-P($A_4^{(1)}$) *(2.6) admits the hypergeometric solution*

$$\widetilde{X}=\frac{F}{\widetilde{F}}, \tag{3.12}$$

$$F=A\,(pa_2{}^{-1}t;p)_\infty\ {}_2\varphi_1\left(\begin{matrix}a_2{}^2,0\\ -p\end{matrix};p,pa_2{}^{-1}t\right)+B\,(pa_2{}^{-1}t;p)_\infty\,(-1)^{\log t/\log p}{}_2\varphi_1\left(\begin{matrix}-a_2{}^2,0\\ -p\end{matrix};p,pa_2{}^{-1}t\right), \tag{3.13}$$

with

$$a_1=p^{-1}a_2{}^2. \tag{3.14}$$

The discrete Riccati equation and the linearized three-term relation are given by

$$\widetilde{X}=\frac{1-pa_2{}^{-1}t}{X+a_2t}, \tag{3.15}$$

and

$$(pa_2{}^{-1}t-1)\widetilde{F}+a_2tF+\utilde{F}=0, \tag{3.16}$$

respectively.

3.3 Symmetric q-P($D_5^{(1)}$)

For convenience, we set

$$X=-iW,\quad a_1=ip^{\nu_1/2},\quad a_2=ip^{\nu_2/2}. \tag{3.17}$$

Symmetric q-P($D_5^{(1)}$) (2.9) can be rewritten as

$$\widetilde{W}\utilde{W}=-\frac{(W+p^{\nu_{1/2}}\tilde{t})(W-p^{-\nu_1/2}\tilde{t})}{(W+p^{\nu_2/2})(W-p^{-\nu_2/2})}. \tag{3.18}$$

Proposition 3.3 ([17, 19]) *Symmetric* q-P($D_5^{(1)}$) *(3.18) admits the hypergeometric solution*

$$W = \frac{\widetilde{G}}{G} - p^{\nu_2/2}, \tag{3.19}$$

$$G = A\, J_{\nu_2}^{(1)}(2ip^{3/4}t^{1/2}; p) + B\, J_{-\nu_2}^{(1)}(2ip^{3/4}t^{1/2}; p), \tag{3.20}$$

with

$$\nu_1 = \nu_2 + 1. \tag{3.21}$$

Here, $J_\nu^{(1)}(x; q)$ *is the Jackson's q-Bessel fuction [5]*

$$J_\nu^{(1)}(x; q) = \frac{(q^{\nu+1}; q)_\infty}{(q; q)_\infty} \left(\frac{x}{2}\right)^\nu {}_2\varphi_1\left(\begin{matrix} 0, 0 \\ q^{\nu+1} \end{matrix}; q, -\frac{x^2}{4}\right). \tag{3.22}$$

The discrete Riccati equation and the linearized three-term relation are given by

$$\widetilde{W} = p^{-\nu_2/2} \frac{W + p^{(\nu_2+1)/2}\tilde{t}}{W + p^{\nu_2/2}}, \tag{3.23}$$

and

$$\widetilde{\widetilde{G}} - (p^{\nu_2/2} + p^{-\nu_2/2})\widetilde{G} + (1 - p^{1/2}\tilde{t})G = 0, \tag{3.24}$$

respectively.

3.4 Symmetric q-P($E_6^{(1)}$)

Proposition 3.4 *Symmetric* q-P($E_6^{(1)}$) *(2.13) admits the hypergeometric solution*

$$X = ({b_1}^{-1} b_5 t - 1)\frac{\widetilde{G}}{G} + b_5 t, \tag{3.25}$$

$$G = A\, {}_2\varphi_1\left(\begin{matrix} {b_1}^{-1}{b_5}^2, -{b_1}^{-1} \\ -p \end{matrix}; p, b_1{b_5}^{-1}\tilde{t}\right) + B\frac{\Theta(t; p)}{\Theta(-t; p)}\, {}_2\varphi_1\left(\begin{matrix} -{b_1}^{-1}{b_5}^2, {b_1}^{-1} \\ -p \end{matrix}; p, b_1{b_5}^{-1}\tilde{t}\right), \tag{3.26}$$

with

$${b_5}^2 = pb_1b_3. \tag{3.27}$$

The discrete Riccati equation and the linearized three-term relation are given by

$$\widetilde{X} = \frac{b_1\tilde{t}X + b_1b_5 - (p{b_1}^2 + {b_5}^2)t}{b_1b_5(X - b_5t)}, \tag{3.28}$$

$$(b_5t - b_1)b_5\widetilde{G} + b_1({b_5}^2 - 1)tG - b_1(b_1t - b_5)\underset{\sim}{G} = 0, \tag{3.29}$$

respectively.

3.5 Symmetric q-P($E_7^{(1)}$)

Proposition 3.5 *Symmetric q-P($E_7^{(1)}$) (2.17) admits the hypergeometric solution*

$$\frac{X - b_1{}^{-1}\tilde{t}}{X - b_5} = \frac{(b_3 b_5 \tilde{t} - 1)}{(b_5^2 - 1)(\tilde{t} - b_3 b_5)} \left((b_5 \tilde{t} - b_1{}^{-1}) \frac{\widetilde{G}}{G} + \tilde{t} - b_1{}^{-1} b_5 \right), \tag{3.30}$$

$$\begin{aligned}
G &= AG_1 + BG_2,\\
G_1 &= \frac{(b_1 b_5 \tilde{t}, -b_3 \tilde{\tilde{t}}; p)_\infty (b_1 b_5 b_3)^{\log t/\log p}}{(b_1{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}, -b_3{}^{-1} \tilde{t}; p)_\infty}\\
&\quad \times {}_8W_7 \left(-b_3{}^2 b_5; p^{1/2} b_3, -p^{1/2} b_3, -b_1 b_3 b_5, b_3 b_5 \tilde{t}, b_3 b_5 \tilde{t}^{-1}; p, p b_1{}^{-1} b_3{}^{-1} b_5{}^{-1} \right), \qquad (3.31)\\
G_2 &= \frac{(\tilde{\tilde{t}}\tilde{t}, -p^{1/2} b_1{}^{-1} b_3{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}, p^{1/2} b_1{}^{-1} b_3{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}, b_3{}^{-1} \tilde{\tilde{t}}, b_1{}^{-1} \tilde{\tilde{t}}; p)_\infty (-b_5)^{\log t/\log p}}{(b_1{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}, p^{1/2} \tilde{t}, -p^{1/2} \tilde{t}, b_3{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}, b_1{}^{-1} b_3{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}^2; p)_\infty}\\
&\quad \times {}_8W_7 \left(b_1{}^{-1} b_3{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}\tilde{t}; b_1{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}, p^{1/2} \tilde{t}, -p^{1/2} \tilde{t}, b_3{}^{-1} b_5{}^{-1} \tilde{\tilde{t}}, p b_1{}^{-1} b_3{}^{-1} b_5{}^{-1}; p, -b_5 \right),
\end{aligned}$$

with

$$b_1 b_3 b_5 b_7 = 1. \tag{3.32}$$

The discrete Riccati equation and the linearized three-term relation are given by

$$\frac{\widetilde{X}X - \tilde{t}^2}{\widetilde{X}X - 1} = \frac{(X - b_1{}^{-1}\tilde{t})(X - b_3{}^{-1}\tilde{t})}{(X - b_5)(X - b_1{}^{-1} b_3{}^{-1} b_5{}^{-1})}, \tag{3.33}$$

$$\frac{(b_1 b_5 \tilde{t} - 1)(b_3 b_5 \tilde{t} - 1)}{(b_1 b_3 b_5 - 1)(1 + b_5)} (\widetilde{G} - G) + (\tilde{t}^2 - 1) G + \frac{(\tilde{t} - b_1 b_5)(\tilde{t} - b_3 b_5)}{(b_1 b_3 b_5 - 1)(1 + b_5)} (G - \underset{\sim}{G}) = 0, \tag{3.34}$$

respectively. To obtain the solution of (3.34), the following proposition is useful:

Proposition 3.6 ([10]) *The three-term relation for $G = AG_1 + BG_2$, where*

$$\begin{aligned}
G_1 =& \frac{(az^{-1})^{\log u/\log p} (abu, acu, adu, bcdz^{-1}u)_\infty}{(bcu, bdu, cdu, azu)_\infty}\\
& \times {}_8W_7(p^{-1} bcdz^{-1}; bz^{-1}, cz^{-1}, dz^{-1}, p^{-1} abcdu, u^{-1}; p, pa^{-1}z), \qquad (3.35)\\
G_2 =& \frac{(az)^{\log u/\log p} (abcdu^2, pbzu, pczu, pdzu, bcdzu)_\infty}{(bcu, bdu, cdu, pu, pbcdzu^2)_\infty}\\
& \times {}_8W_7(bcdzu^2; bcu, bdu, cdu, pu, pa^{-1}z; p, az), \qquad (3.36)
\end{aligned}$$

is given by

$$\begin{aligned}
&\frac{(1 - abu)(1 - acu)(1 - adu)(1 - p^{-1}abcdu)}{a(1 - p^{-1}abcdu^2)(1 - abcdu^2)} (G(pu) - G(u)) + (a + a^{-1} - z - z^{-1}) G(u)\\
&- \frac{a(1 - p^{-1}bcu)(1 - p^{-1}bdu)(1 - p^{-1}cdu)(1 - u)}{(1 - p^{-2}abcdu^2)(1 - p^{-1}abcdu^2)} (G(u) - G(p^{-1}u)) = 0.
\end{aligned} \tag{3.37}$$

Here A and B are quasi-constants with respect to u.

Substituting

$$\begin{aligned} &a = ib_1{}^{1/2}b_3{}^{1/2}b_5, \quad b = ip^{1/2}b_1{}^{-1/2}b_3{}^{1/2}, \quad c = -ip^{1/2}b_1{}^{-1/2}b_3{}^{1/2}, \\ &d = -ib_1{}^{1/2}b_3{}^{1/2}b_5, \quad z = ib_1{}^{-1/2}b_3{}^{-1/2}, \quad u = b_3{}^{-1}b_5{}^{-1}t, \end{aligned} \tag{3.38}$$

in Proposition 3.6, we obtain the solution of (3.34).

For later convenience, we rewrite the solution in the following manner:

$$F = \frac{(b_1{}^{-1}b_5{}^{-1}\tilde{\tilde{t}}, -b_3{}^{-1}\tilde{t}; p)_\infty}{(b_1b_5\tilde{t}, -b_3\tilde{\tilde{t}}; p)_\infty (b_1b_5b_3)^{\log t/\log p}} G. \tag{3.39}$$

Then Proposition 3.5 is rephrased as follows:

Proposition 3.7 *Symmetric* q-P$(E_7^{(1)})$ *(2.17) admits the hypergeometric solution*

$$\frac{X - b_1{}^{-1}\tilde{t}}{X - b_5} = \frac{b_3b_5\tilde{t} - 1}{b_1(b_5{}^2 - 1)(\tilde{t} - b_3b_5)} \left[\frac{(\tilde{\tilde{t}} - b_1b_5)(\tilde{t} + b_3)}{b_3\tilde{\tilde{t}} + 1} \frac{\widetilde{F}}{F} + b_1\tilde{t} - b_5 \right], \tag{3.40}$$

$$\begin{aligned} F &= AF_1 + BF_2, \\ F_1 &= {}_8W_7\left(-b_3{}^2b_5; p^{1/2}b_3, -p^{1/2}b_3, -b_1b_3b_5, b_3b_5\tilde{t}, b_3b_5\tilde{t}^{-1}; p, pb_1{}^{-1}b_3{}^{-1}b_5{}^{-1}\right), \\ F_2 &= \frac{(-b_3{}^{-1}\tilde{t}, \tilde{\tilde{t}}\tilde{t}, -p^{1/2}b_1{}^{-1}b_3{}^{-1}b_5{}^{-1}\tilde{\tilde{t}}, p^{1/2}b_1{}^{-1}b_3{}^{-1}b_5{}^{-1}\tilde{\tilde{t}}, b_3{}^{-1}\tilde{\tilde{t}}, b_1{}^{-1}\tilde{\tilde{t}}; p)_\infty}{(b_1b_5\tilde{t}, -b_3\tilde{\tilde{t}}, p^{1/2}\tilde{t}, -p^{1/2}\tilde{t}, b_3{}^{-1}b_5{}^{-1}\tilde{\tilde{t}}, b_1{}^{-1}b_3{}^{-1}b_5{}^{-1}\tilde{\tilde{t}}^2; p)_\infty(-b_1b_3)^{\log t/\log p}} \\ &\quad \times {}_8W_7\left(b_1{}^{-1}b_3{}^{-1}b_5{}^{-1}\tilde{\tilde{t}}\tilde{t}; b_1{}^{-1}b_5{}^{-1}\tilde{\tilde{t}}, p^{1/2}\tilde{t}, -p^{1/2}\tilde{t}, b_3{}^{-1}b_5{}^{-1}\tilde{\tilde{t}}, pb_1{}^{-1}b_3{}^{-1}b_5{}^{-1}; p, -b_5\right). \end{aligned} \tag{3.41}$$

when

$$b_1b_3b_5b_7 = 1. \tag{3.42}$$

We note that the linear three-term relation for F is given by

$$\begin{aligned} &\frac{(b_1b_5\tilde{t} - 1)(b_3b_5\tilde{t} - 1)}{(b_1b_3b_5 - 1)(1 + b_5)} \left(\frac{(b_1b_5 - \tilde{\tilde{t}})(b_3 + \tilde{t})}{(1 - b_1b_5\tilde{t})(1 + b_3\tilde{\tilde{t}})} \widetilde{F} - F \right) + (\tilde{t}^2 - 1)F \\ &\quad + \frac{(\tilde{t} - b_1b_5)(\tilde{t} - b_3b_5)}{(b_1b_3b_5 - 1)(1 + b_5)} \left(F - \frac{(1 - b_1b_5t)(1 + b_3\tilde{t})}{(b_1b_5 - \tilde{t})(b_3 + t)} \underset{\sim}{F} \right) = 0. \end{aligned} \tag{3.43}$$

3.6 Symmetric q-P$(E_8^{(1)})$

Proposition 3.8 *Symmetric* q-P$(E_8^{(1)})$ *(2.21) admits the hypergeometric solution*

$$\frac{X - \beta_3}{X - \beta_1} = \frac{(\alpha_3 - \alpha_5)(\beta_1 - \beta_5)}{(\alpha_1 - \alpha_5)(\beta_3 - \beta_5)} \left[(\alpha_1 - \alpha_3) \frac{\widetilde{G}}{G} + \alpha_3 - \tilde{\beta}_1 \right], \tag{3.44}$$

$$
\begin{aligned}
G &= AG_1 + BG_2,\\
G_1 &= \frac{(b_1b_3\tilde{t}^2, b_1b_5\tilde{t}^2, -\tilde{t}^2, {b_1}^{-1}{b_3}^{-1}{b_5}^{-1}\tilde{t}^2; p^2)_\infty}{(-\tilde{t}^2, {b_1}^{-1}{b_5}^{-1}\tilde{t}^2, {b_1}^{-1}{b_3}^{-1}\tilde{t}^2, b_1b_3b_5\tilde{t}^2; p^2)_\infty}(-b_1)^{\log t/\log p}\\
&\quad \times {}_8W_7(p^{-2}{b_1}^{-1}{b_3}^{-2}{b_5}^{-2}; {b_5}^{-1}, {b_3}^{-1}, -{b_1}^{-1}{b_3}^{-1}{b_5}^{-1}, {b_3}^{-1}{b_5}^{-1}t^2, {b_3}^{-1}{b_5}^{-1}\tilde{t}^{-2}; p^2, -p^2{b_1}^{-1}), \qquad (3.45)\\
G_2 &= \frac{(\tilde{t}^4, -b_3\tilde{\tilde{t}}^2, -b_5\tilde{\tilde{t}}^2, {b_1}^{-1}\tilde{\tilde{t}}^2, -{b_1}^{-1}{b_3}^{-1}{b_5}^{-1}\tilde{t}^2; p^2)_\infty}{(-\tilde{t}^2, {b_1}^{-1}{b_5}^{-1}\tilde{t}^2, {b_1}^{-1}{b_3}^{-1}\tilde{t}^2, b_3b_5\tilde{\tilde{t}}^2, -{b_1}^{-1}\tilde{\tilde{t}}^2\tilde{t}^2; p^2)_\infty}{b_1}^{\log t/\log p}\\
&\quad \times {}_8W_7(-{b_1}^{-1}\tilde{t}^4; -\tilde{t}^2, {b_1}^{-1}{b_5}^{-1}\tilde{t}^2, {b_1}^{-1}{b_3}^{-1}\tilde{t}^2, b_3b_5\tilde{\tilde{t}}^2, -p^2{b_1}^{-1}; p^2, b_1),
\end{aligned}
$$

with

$$b_1b_3b_5b_7 = p^{-2}. \tag{3.46}$$

Here, α_i and β_i are given by

$$\alpha_i = b_i\tilde{t} + \frac{1}{b_i\tilde{t}}, \quad \beta_i = \frac{t}{b_i} + \frac{b_i}{t}. \tag{3.47}$$

The discrete Riccati equation and the linearized three-term relation are given by

$$\widetilde{X} = \frac{X\begin{vmatrix}1 & \alpha_1 & \alpha_1\beta_1\\ 1 & \alpha_3 & \alpha_3\beta_3\\ 1 & \alpha_5 & \alpha_5\beta_5\end{vmatrix} + \begin{vmatrix}\alpha_1 & \beta_1 & \alpha_1\beta_1\\ \alpha_3 & \beta_3 & \alpha_3\beta_3\\ \alpha_5 & \beta_5 & \alpha_5\beta_5\end{vmatrix}}{X\begin{vmatrix}1 & \alpha_1 & \beta_1\\ 1 & \alpha_3 & \beta_3\\ 1 & \alpha_5 & \beta_5\end{vmatrix} + \begin{vmatrix}1 & \beta_1 & \alpha_1\beta_1\\ 1 & \beta_3 & \alpha_3\beta_3\\ 1 & \beta_5 & \alpha_5\beta_5\end{vmatrix}}, \tag{3.48}$$

$$
\begin{aligned}
&\frac{(b_1b_3\tilde{t}^2-1)(b_1b_5\tilde{t}^2-1)(t^2-b_3b_5)}{(\tilde{t}^2-1)(\tilde{t}^2t^2-1)}(\widetilde{G}-G) + b_3b_5({b_1}^2-1)G\\
&\quad + \frac{(t^2-b_1b_3)(t^2-b_1b_5)(b_3b_5\tilde{t}^2-1)}{(t^2-1)(\tilde{t}^2t^2-1)}(G-\underset{\sim}{G}) = 0,
\end{aligned} \tag{3.49}
$$

respectively. We note that (3.45) is obtained by substituting

$$a = -ib_1, \quad b = i{b_5}^{-1}, \quad c = i{b_3}^{-1}, \quad d = -i{b_1}^{-1}{b_3}^{-1}{b_5}^{-1}, \quad z = i, \quad u = b_3b_5\tilde{t}^2, \tag{3.50}$$

in Proposition 3.6.

4 Continuous limits

In this section, we discuss the continuous limits of the symmetric q-Painlevé equations and their hypergeometric solutions. We introduce the hypergeometric series ${}_sF_r$ defined by

$${}_sF_r\left(\begin{matrix}a_1, \cdots, a_s\\ b_1, \cdots, b_r\end{matrix}; z\right) = \sum_{n=0}^{\infty} \frac{(a_1)_n \cdots (a_s)_n}{(b_1)_n \cdots (b_r)_n n!} z^n, \tag{4.1}$$

where $(a)_n = a(a+1)\ldots(a+n-1)$.

4.1 Symmetric q-P$((A_2+A_1)^{(1)})$

It is known that symmetric q-P$((A_2+A_1)^{(1)})$ (2.4) yields the Painlevé II equation

$$z'' = 2z^3 + sz + \alpha, \tag{4.2}$$

where $z' = \mathrm{d}z/\mathrm{d}s$, by putting

$$t = -2e^{-s\epsilon^2/4-\alpha\epsilon^3/4}, \quad c = e^{(1+2\alpha)\epsilon^3/4}, \quad X = e^{-s\epsilon^2/4+\alpha\epsilon^3/4}(1+\epsilon z), \quad p = e^{-\epsilon^3/4}, \tag{4.3}$$

and taking the limit $\epsilon \to +0$ [37]. The limit of the hypergeometric solution is given as follows:

Proposition 4.1 *Under the parametrization (4.3), the hypergeometric solution in Proposition 3.1 is reduced to the following solution to (4.2):*

$$z = \frac{G'}{G}, \tag{4.4}$$

$$G = A\,\mathrm{Ai}(2^{-1/3}se^{-\pi i/3}) + B\,\mathrm{Ai}(2^{-1/3}se^{\pi i/3}), \tag{4.5}$$

with

$$\alpha = -\frac{1}{2}. \tag{4.6}$$

Here A and B are constants, and $\mathrm{Ai}(x)$ *is the Airy function*

$$\mathrm{Ai}(x) = \frac{1}{2\pi i}\int_{-\infty}^{\infty} e^{iu^3/3+ixu}du. \tag{4.7}$$

We note that (3.9) and (3.10) are reduced to the following equations

$$z' = -z^2 - \frac{s}{2}, \tag{4.8}$$

$$G'' + \frac{s}{2}G = 0, \tag{4.9}$$

respectively. We also note that (4.5) is obtained from (3.7) by using the following proposition:

Proposition 4.2 ([9]) *With the substitutions*

$$q = e^{-\delta^3/2}, \quad T = -2ie^{-(u/2)\delta^2}, \tag{4.10}$$

as $\delta \to +0$, it follows that

$${}_1\varphi_1\left(\begin{matrix}0\\-q\end{matrix};q,qT\right) = 2\pi^{1/2}\delta^{-1/2}e^{\log 2(\pi i/\delta^3)-(\pi i/2\delta)u-\pi i/12}\left[\mathrm{Ai}(ue^{-\pi i/3}) + O(\delta^2)\right], \tag{4.11}$$

$${}_1\varphi_1\left(\begin{matrix}0\\-q\end{matrix};q,-qT\right) = 2\pi^{1/2}\delta^{-1/2}e^{-\log 2(\pi i/\delta^3)+(\pi i/2\delta)u+\pi i/12}\left[\mathrm{Ai}(ue^{\pi i/3}) + O(\delta^2)\right], \tag{4.12}$$

for u in any compact domain of $\mathbb{C}$.

4.2 Symmetric q-P$(A_4^{(1)})$

Putting

$$t = 2ie^{is\epsilon - 3\alpha\epsilon^2/2}, \quad a_1 = -e^{-2^{-1/2}i\beta^{1/2}\epsilon^2}, \quad a_2 = ie^{-\alpha\epsilon^2/2}, \quad X = 1 - i\epsilon z, \quad p = e^{-\epsilon^2}, \tag{4.13}$$

and taking the limit $\epsilon \to 0$, symmetric q-P$(A_4^{(1)})$ (2.6) yields the Painlevé IV equation [37]

$$z'' = \frac{(z')^2}{2z} + \frac{3}{2}z^3 + 4sz^2 + 2(s^2 - \alpha)z + \frac{\beta}{z}. \tag{4.14}$$

Proposition 4.3 *Under the parametrization (4.13), the hypergeometric solution in Proposition 3.2 is reduced to*

$$z = \frac{F'}{F}, \tag{4.15}$$

$$F = A\, e^{-s^2} D_{-\alpha}(2^{1/2}s) + B\, D_{\alpha-1}(2^{1/2}is), \quad s \notin \mathbb{R},\ \sqrt{-1}\mathbb{R}, \tag{4.16}$$

with

$$\alpha = 1 + 2^{-1/2}i\beta^{1/2}. \tag{4.17}$$

Here A and B are constants, and $D_\lambda(x)$ is the Weber function

$$D_\lambda(x) = \int_\gamma e^{-u^2/2 - xu} u^{-\lambda-1} du, \tag{4.18}$$

where the contour of integration (denote it by γ) runs from $-\infty$ to $+\infty$ so that $u = 0$ lie to the right of the contour.

Remark 4.4 *The hypergeometric solution to the Painlevé IV equation given in Proposition 4.3 is consistent with the solution appeared in [28,31].*

Note that (3.15) and (3.16) are reduced to the following equations

$$z' = -z^2 - 2sz + 2(\alpha - 1), \tag{4.19}$$

$$F'' + 2sF' + 2(1 - \alpha)F = 0. \tag{4.20}$$

respectively. We also note that (4.16) is obtained from (3.13) by using the following lemmas:

Lemma 4.5 *With the substitutions (4.13), it follows that*

$$\begin{aligned} &{}_2\varphi_1\left(\begin{matrix} a_2{}^2, 0 \\ -p \end{matrix}; p, pa_2^{-1}t\right) - \frac{(a_2{}^{-2}; p)_\infty e^{\pi i \log(pa_2{}^{-1})/\log p}}{(-a_2{}^2; p)_\infty (-1)^{\log t/\log p}}\, {}_2\varphi_1\left(\begin{matrix} -a_2{}^2, 0 \\ -p \end{matrix}; p, pa_2{}^{-1}t\right) \\ &= \frac{(-1)^{3\alpha/2} 2^{-\alpha/2} e^{(\pi i \log 2 + \pi^2/4)/\epsilon^2 - \pi s/\epsilon}}{\pi^{1/2}\epsilon^{1-\alpha}} \int_L e^{-u^2/2 - 2^{1/2}su} u^{\alpha-1} [1 + O(\epsilon)]\, du, \end{aligned} \tag{4.21}$$

as

$$\epsilon \to +0 \quad (\text{when } \mathrm{Im}(s) < 0), \tag{4.22}$$

$$\epsilon \to -0 \quad (\text{when } \mathrm{Im}(s) > 0). \tag{4.23}$$

The contour L runs from $-\infty$ to $+\infty$ when $\mathrm{Im}(s) < 0$ and $+\infty$ to $-\infty$ when $\mathrm{Im}(s) > 0$ so that $u = 0$ lie to the right of the contour.

Lemma 4.6 *With the substitutions (4.13), it follows that*

$$
\begin{aligned}
&{}_2\varphi_1\left(\begin{matrix}-a_2{}^2, 0\\ -p\end{matrix}; p, pa_2{}^{-1}t\right)\\
&\quad - \frac{(-a_2{}^2;p)_\infty e^{(2\log(-pa_2)\log a_2 - \log(pa_2{}^{-1})\log(-a_2{}^{-2}))/\log p}}{a_2{}^4(a_2{}^2;p)_\infty}(-1)^{\log t/\log p}{}_2\varphi_1\left(\begin{matrix}a_2{}^2, 0\\ -p\end{matrix}; p, pa_2{}^{-1}t\right)\\
&= (-1)^{-1-4\alpha}2^{-1/2+\alpha/2}\pi^{-1/2}\epsilon^{-\alpha}e^{s^2+2\pi i\log 2/\epsilon^2}\int_{\hat{L}_2} e^{-u^2/2-2^{1/2}isu}u^{-\alpha}[1+O(\epsilon)]du,
\end{aligned}
$$

as

$$\epsilon \to +0 \quad (\text{when } \mathrm{Re}(s) < 0), \tag{4.24}$$

$$\epsilon \to -0 \quad (\text{when } \mathrm{Re}(s) > 0). \tag{4.25}$$

The contour L runs from $-\infty$ to $+\infty$ when $\mathrm{Re}(s) > 0$ and $+\infty$ to $-\infty$ when $\mathrm{Re}(s) < 0$ so that $u = 0$ lie to the right of the contour.

The proof of Lemma 4.5 and 4.6 will be given in the next section. Moreover, for the asymptotic expansions of the q-shifted factorials as $q \to 1^-$, we use the following proposition:

Proposition 4.7 ([23, 34]) *As $q \to 1^-$, the q-shifted factorials have an asymptotic expansions,*

$$\log(t;q)_\infty = \frac{Li_2(t)}{\log q} + \frac{\log(1-t)}{2} + O(\log q), \tag{4.26}$$

$$\log(q;q)_\infty = \frac{\pi^2}{6\log q} + \frac{1}{2}\log\left(-\frac{2\pi}{\log q}\right) + O(\log q), \tag{4.27}$$

$$\log(-q;q)_\infty = -\frac{\pi^2}{12\log q} - \frac{\log 2}{2} + O(\log q). \tag{4.28}$$

This is uniform for $t \in \mathbb{C}$ such that $|\arg(1-t)| < \pi$. Here $Li_2(t)$ is the Euler dilogarithm defined by

$$Li_2(t) = -\int_0^t \frac{\log(1-u)}{u}du \tag{4.29}$$

$$= \sum_{k=1}^{\infty}\frac{t^k}{k^2} \quad (|t| < 1). \tag{4.30}$$

4.3 Symmetric q-P$(D_5^{(1)})$

Putting

$$t^{1/2} = \frac{i(1-p)}{2}s, \quad W = \frac{\epsilon s}{2}z, \quad p = 1+\epsilon, \tag{4.31}$$

and taking the limit $\epsilon \to -0$, symmetric q-P$(D_5^{(1)})$ (3.18) yields the Painlevé III equation [38]

$$z'' - \frac{(z')^2}{z} + \frac{z'}{s} + \frac{2(\nu_1 - \nu_2 z^2)}{s}) - z^3 + \frac{1}{z} = 0. \tag{4.32}$$

Proposition 4.8 ([17, 19]) *Under the parametrization (4.31), the hypergeometric solution in Proposition 3.3 is reduced to*

$$z = \frac{G'}{G} - \frac{\nu_2}{s}, \tag{4.33}$$

$$G = A\, J_{\nu_2}(s) + B\, J_{-\nu_2}(s), \tag{4.34}$$

with

$$\nu_1 = \nu_2 + 1. \tag{4.35}$$

Here, A and B are constants and $J_\nu(x)$ is the Bessel function

$$J_\nu(x) = \frac{x^\nu}{2^\nu \Gamma(\nu+1)}\, {}_0F_1\left(\begin{matrix} - \\ \nu+1 \end{matrix}; -\frac{x^2}{4}\right). \tag{4.36}$$

We note that (3.23) and (3.24) are reduced to the following equations

$$z' + z^2 + \frac{1+2\nu_2}{s} z + 1 = 0, \tag{4.37}$$

$$G'' + \frac{G'}{s} + \left(1 - \frac{{\nu_2}^2}{s^2}\right) G = 0, \tag{4.38}$$

respectively.

4.4 Symmetric q-P$(E_6^{(1)})$

Putting

$$\begin{aligned} &t = \frac{(\gamma^2\epsilon^2 - 16)^{1/2}}{4\epsilon}(1+\epsilon)^{-\log s/\epsilon}, \quad b_1 = -1 + 2^{1/2}\beta\epsilon, \quad b_3 = 1 + 2^{1/2}\alpha\epsilon, \\ &b_5 = \frac{(\gamma^2\epsilon^2 - 16)^{1/2}}{\gamma\epsilon - 4}, \quad X = \frac{z+1}{z-1}, \quad p = 1 + \epsilon, \end{aligned} \tag{4.39}$$

and taking the limit $\epsilon \to -0$, symmetric q-P$(E_6^{(1)})$ (2.13) yields the Painlevé V equation [37]

$$z'' = \left(\frac{1}{2z} + \frac{1}{z-1}\right)(z')^2 - \frac{z'}{s} + \frac{(z-1)^2}{s^2}\left(\alpha^2 z - \frac{\beta^2}{z}\right) + \frac{\gamma}{s} z - \frac{2z(z+1)}{z-1}. \tag{4.40}$$

Proposition 4.9 *Under the parametrization (4.39), the hypergeometric solution in Proposition 3.4 is reduced to the following solution to (4.40):*

$$z = \frac{sG' - 2^{1/2}\beta G}{sG' - (2^{1/2}\beta + 2s)G}. \tag{4.41}$$

$$G = A\, {}_2F_0\left(\begin{matrix} 2^{1/2}\beta + \frac{\gamma}{2}, 2^{1/2}\beta \\ - \end{matrix}; -\frac{1}{2s}\right) + \hat{B} e^{2s} s^{2^{-1}\gamma + 2^{3/2}\beta - 1}\, {}_2F_0\left(\begin{matrix} 1 - 2^{1/2}\beta - \frac{\gamma}{2}, 1 - 2^{1/2}\beta \\ - \end{matrix}; \frac{1}{2s}\right), \tag{4.42}$$

with

$$\gamma = 2(1 + 2^{1/2}\alpha - 2^{1/2}\beta). \tag{4.43}$$

Here, A and $\hat{B}$ are constants.

The hypergeometric solution in Proposition 3.4 is reduced to the following solution to (4.40)

$$sz' + \left(2^{1/2}\beta + \frac{\gamma}{2} - 1\right)z^2 - \left(2^{3/2}\beta + \frac{\gamma}{2} + 2s - 1\right)z + 2^{1/2}\beta = 0, \tag{4.44}$$

$$G'' + \frac{2 - \gamma - 2^{5/2}\beta - 4s}{2s}G' + \frac{\beta\gamma + 2^{3/2}\beta^2}{2^{1/2}s^2}G = 0, \tag{4.45}$$

respectively. We note that (4.42) is obtained from (3.26) by the following procedure. By using the Heine's transformation [20]

$${}_2\varphi_1\left(\begin{matrix}a, b\\ c\end{matrix}; q, z\right) = \frac{(abc^{-1}, z; q)_\infty}{(z; q)_\infty}\,{}_2\varphi_1\left(\begin{matrix}a^{-1}c, b^{-1}c\\ c\end{matrix}; q, abc^{-1}z\right), \tag{4.46}$$

and setting

$$\hat{B} = \frac{\Theta(t; p)\Theta(-b_1{b_5}^{-1}\epsilon\tilde{t}; p)({b_1}^{-1}b_5t, pb_1{b_5}^{-1}t^{-1}; p)_\infty}{\Theta(-t; p)\Theta({b_1}^{-1}b_5\epsilon t; p)(b_1{b_5}^{-1}\tilde{t}, {b_1}^{-1}b_5t^{-1}; p)_\infty}B, \tag{4.47}$$

(3.26) is rewritten as

$$\begin{aligned} G =& A\,{}_2\varphi_1\left(\begin{matrix}{b_1}^{-1}{b_5}^2, -{b_1}^{-1}\\ -p\end{matrix}; p, b_1{b_5}^{-1}\tilde{t}\right)\\ &+ \hat{B}\frac{\Theta({b_1}^{-1}b_5\epsilon t; p)({b_1}^{-1}b_5t^{-1}; p)_\infty}{\Theta(-b_1{b_5}^{-1}\epsilon\tilde{t}; p)(pb_1{b_5}^{-1}t^{-1}; p)_\infty}\,{}_2\varphi_1\left(\begin{matrix}pb_1{b_5}^{-2}, -pb_1\\ -p\end{matrix}; p, {b_1}^{-1}b_5t\right). \end{aligned} \tag{4.48}$$

Then (4.48) yields the hypergeometric series in (4.42) by taking a term-by-term limit. We use Proposition 4.7 for the limit of the coefficient.

Remark 4.10 *Setting*

$$s = -\frac{t}{2}, \quad z(s) = 1 - \frac{1}{f(t)}, \quad G(s) = e^{2s}s^{a_2}\phi(t), \quad \alpha = -2^{-1/2}a_1, \quad \beta = 2^{-1/2}a_2, \tag{4.49}$$

we can rewrite the Riccati equation, the dependent variable teansformation and the linear differential equation as

$$tf' - tf(1 - f) + (a_1 + a_2)f - a_1 = 0, \tag{4.50}$$

$$f = \frac{\phi'}{\phi}, \tag{4.51}$$

$$t\phi'' + (a_1 + a_2 - t)\phi' - a_1\phi = 0, \tag{4.52}$$

respectively. These coincide with the result in [22,32]. We note here that in [22,32] the solution to (4.52) is given by the hypergeometric series around zero while in this section that is given by the series at infinity.

4.5 Symmetric q-P($E_7^{(1)}$)

Putting

$$b_1 = e^{(\beta+1/2)\epsilon}, \quad b_3 = -e^{(\alpha+1/2)\epsilon}, \quad b_5 = e^{\delta\epsilon}, \quad b_7 = -e^{\gamma\epsilon}, \quad p = e^{\epsilon}, \tag{4.53}$$

and taking the limit $\epsilon \to -0$, symmetric q-P($E_7^{(1)}$) (2.17) yields Painlevé VI equation

$$\begin{aligned} X'' =& \frac{1}{2}\left(\frac{1}{X+1} + \frac{1}{X-1} + \frac{1}{X+t} + \frac{1}{X-t}\right)(X')^2 \\ & - \left(\frac{1}{t} + \frac{1}{t-1} + \frac{1}{t+1} + \frac{1}{X-t} - \frac{1}{X+t}\right)X' \\ & + \frac{(X^2-t^2)(X^2-1)}{t^2(t^2-1)}\left(\frac{(4\alpha^2-1)t}{4(X+t)^2} - \frac{(4\beta^2-1)t}{4(X-t)^2} - \frac{\gamma^2}{(X+1)^2} + \frac{\delta^2}{(X-1)^2}\right), \end{aligned} \tag{4.54}$$

where $X' = \mathrm{d}X/\mathrm{d}t$ [6].

Proposition 4.11 *Under the parametrization (4.53), the hypergeometric solution in Proposition 3.5 is reduced to the following solution to (4.54):*

$$X(t) = \frac{(t-1)tF' - (1+\alpha+\beta+t(\alpha+\beta+2\delta+1))F}{(t-1)tF' - (1+\alpha+\beta+2\delta+t(\alpha+\beta+1))F}. \tag{4.55}$$

$$\begin{aligned} F =& A\ {}_2F_1\left(\begin{matrix}\alpha+1, \alpha+\beta+\delta+1 \\ \alpha+\delta+1\end{matrix}; \left(\frac{1+t}{1-t}\right)^2\right) \\ & + \hat{B}\ \frac{(1-t)^{2(1+2\alpha+\beta+\delta)}}{(1+t)^{2(\alpha+\beta)}t^{1+\alpha+\beta}}\ {}_2F_1\left(\begin{matrix}-\alpha-\beta-\delta, -\alpha \\ -\alpha-\beta\end{matrix}; -\frac{4t}{(1-t)^2}\right), \end{aligned} \tag{4.56}$$

with

$$\alpha+\beta+\delta+\gamma+1 = 0. \tag{4.57}$$

Here, A and $\hat{B}$ are constants.

We note that (3.33) and (3.43) are reduced to the following equations

$$X' = \frac{1+\alpha+\beta+2\delta-t(\alpha-\beta)}{t(t^2-1)}X^2 + \frac{\alpha+\beta+1}{t}X + \frac{\alpha-\beta-t(\alpha+\beta+2\delta+1)}{t^2-1}, \tag{4.58}$$

$$F'' - \frac{2+\alpha+\beta+2(2+3\alpha+\beta+2\delta)t+(\alpha+\beta)t^2}{t(t^2-1)}F' + \frac{4(\alpha+1)(\alpha+\beta+\delta+1)}{t(t-1)^2}F = 0, \tag{4.59}$$

respectively. We also note that (4.56) is obtained by the following procedure. By using the transformation (see, (III.23) in [5])

$${}_8W_7\left(a; b, c, d, e, f; q, \frac{q^2a^2}{bcdef}\right) = \frac{\left(qa, \frac{qa}{ef}, \frac{q^2a^2}{bcde}, \frac{q^2a^2}{bcdf}; q\right)_\infty}{\left(\frac{qa}{e}, \frac{qa}{f}, \frac{q^2a^2}{bcd}, \frac{q^2a^2}{bcdef}; q\right)_\infty}\ {}_8W_7\left(\frac{qa^2}{bcd}; \frac{qa}{cd}, \frac{qa}{bd}, \frac{qa}{bc}, e, f; q, \frac{qa}{ef}\right), \tag{4.60}$$

and setting

$$\hat{B} = B\frac{\Theta(t;p)}{\Theta(-b_1b_3t;p)(-b_1b_3)^{\log t/\log p}}, \tag{4.61}$$

(3.41) is rewritten as

$$\begin{aligned}
&F = AF_1 + \hat{B}\hat{F}_2,\\
&F_1 = {}_8W_7\left(-{b_3}^2b_5; p^{1/2}b_3, -p^{1/2}b_3, -b_1b_3b_5, b_3b_5\tilde{t}, b_3b_5\tilde{t}^{-1}; p, p{b_1}^{-1}{b_3}^{-1}{b_5}^{-1}\right),\\
&\hat{F}_2 = \frac{(-{b_3}^{-1}\tilde{t}, -p^{1/2}{b_1}^{-1}{b_3}^{-1}{b_5}^{-1}\tilde{\tilde{t}}, p^{1/2}{b_1}^{-1}{b_3}^{-1}{b_5}^{-1}\tilde{\tilde{t}}, {b_3}^{-1}\tilde{\tilde{t}}, b_3b_5\tilde{t}, -p{b_1}^{-1}{b_3}^{-1}, -{b_3}^{-1}\tilde{\tilde{t}}; p)_\infty}{(b_1b_5\tilde{t}, -b_3\tilde{\tilde{t}}, p^{1/2}\tilde{t}, -p^{1/2}\tilde{t}, {b_3}^{-1}{b_5}^{-1}\tilde{\tilde{t}}, -p^3{b_1}^{-1}{b_3}^{-2}{b_5}^{-1}t, -b_5; p)_\infty}\\
&\quad\times\frac{\Theta(-b_1b_3t;p)}{\Theta(t;p)}\\
&\quad\times {}_8W_7\left(-{b_1}^{-1}{b_3}^{-2}{b_5}^{-1}\tilde{\tilde{t}}; -p{b_1}^{-1}{b_3}^{-1}{b_5}^{-1}, -p^{1/2}{b_3}^{-1}, p^{1/2}{b_3}^{-1}, {b_3}^{-1}{b_5}^{-1}\tilde{\tilde{t}}, p{b_1}^{-1}{b_3}^{-1}{b_5}^{-1}; p, b_3b_5\tilde{t}\right).
\end{aligned} \tag{4.62}$$

Then (4.56) is obtained from (4.62) by taking a term-by-term limit and using Proposition 4.7.

5 Proof of Lemma 4.5 and 4.6

In this section, we give the proof of Lemma 4.5 and 4.6. Taking the continuous limit term-by-term in the hypergeometric series in (3.13) does not yield any meaningful result. In this case, we use the saddle point method [9, 34, 44]. Consider the complex integral

$$I(s) = \int_C e^{-g(z)/\epsilon^2} f(z)\,dz, \tag{5.1}$$

where $f(z)$ and $g(z)$ are analytic functions. The point $z = z_0$ satisfying $g'(z_0) = 0$ and the direction

$$\arg(z - z_0) = \frac{\pi}{2} - \frac{1}{2}\arg\left(-\frac{g''(z_0)}{\epsilon^2}\right), \tag{5.2}$$

are called a saddle point and a steepest descent direction, respectively. When the path C passes in the steepest descent direction, $|e^{-g(z)/\epsilon^2} f(z)|$ reaches its peak at the saddle point. Therefore $I(s)$ may be evaluated only around the saddle point when $\epsilon \to 0$. For the asymptotic expansions of the q-shifted factorials as $q \to 1^-$, we use Proposition 4.7.

We here fix the branch of log and fractional power functions as

$$\log z = \log|z| + i\arg z \quad (0 \le \arg z < 2\pi), \tag{5.3}$$

$$z^{m/n} = e^{(m/n)\log z} = |z|^{m/n}e^{(m/n)i\arg z} \quad (m \in \mathbb{Z},\ n \in \mathbb{Z}_{>0},\ 0 \le \arg z < 2\pi), \tag{5.4}$$

where $z \in \mathbb{C}^*$. Note that $\log(XY) = \log X + \log Y$ and $\log(X/Y) = \log X - \log Y$ are valid only mod $2\pi i$.

5.1 Proof of Lemma 4.5

We consider the continuous limit of

$$A_1 = {}_2\varphi_1\left(\begin{matrix}{a_2}^2, 0\\ -p\end{matrix}; p, pa_2^{-1}t\right), \tag{5.5}$$

with

$$t = 2ie^{is\epsilon - 3\alpha\epsilon^2/2}, \quad a_2 = ie^{-\alpha\epsilon^2/2}, \quad p = e^{-\epsilon^2}, \quad \epsilon \to \pm 0. \tag{5.6}$$

We note that in taking the limit, the sign of ϵ is chosen according to the value of s as shown later. By using the Heine's transformation [20]

$${}_2\varphi_1\left(\begin{matrix}a, 0\\ c\end{matrix}; q, z\right) = \frac{(az;q)_\infty}{(c,z;q)_\infty}\, {}_1\varphi_1\left(\begin{matrix}z\\ az\end{matrix}; q, c\right), \tag{5.7}$$

A_1 is rewritten as

$$A_1 = \frac{(pa_2t;p)_\infty}{(-p, p{a_2}^{-1}t;p)_\infty}\, {}_1\varphi_1\left(\begin{matrix}p{a_2}^{-1}t\\ pa_2t\end{matrix}; p, -p\right). \tag{5.8}$$

We next prepare a suitable integral representation for A_1.

Lemma 5.1 *It holds that*

$${}_1\varphi_1\left(\begin{matrix}a\\ c\end{matrix}; q, x\right) = \frac{(a;q)_\infty (q;q)_\infty}{2\pi i (c;q)_\infty} \int_C \frac{z^{-\log x/\log q}(cz^{-1};q)_\infty}{(az^{-1};q)_\infty (z;q)_\infty}\, dz, \tag{5.9}$$

where the contour C runs from $-i\infty$ to $i\infty$ so that the poles of $1/(z;q)_\infty$ lie to the right of the contour and the other poles lie to the left of the contour (*as shown in Figure 1*).

Proof. We consider the integral

$$\oint_{C_N^0} \frac{z^{-\log x/\log q}(cz^{-1};q)_\infty}{(az^{-1};q)_\infty (z;q)_\infty}\, dz, \tag{5.10}$$

where the contour C_N^0 is a large clockwise-oriented semicircle of radius $q^{-N-1/2}$ with center at the origin and circle round only part of the poles of $1/(z;q)_\infty$ (as shown in Figure 2). By the residue theorem, we have

$$\frac{1}{2\pi i}\oint_{C_N^0} \frac{z^{-\log x/\log q}(cz^{-1};q)_\infty}{(az^{-1};q)_\infty (z;q)_\infty}\, dz = \frac{(c;q)_\infty}{(a;q)_\infty (q;q)_\infty} \sum_{n=0}^{N} \frac{(a;q)_n}{(c;q)_n (q;q)_n} (-1)^n q^{n(n-1)/2} x^n. \tag{5.11}$$

Therefore we obtain

$${}_1\varphi_1\left(\begin{matrix}a\\ c\end{matrix}; q, x\right) = \frac{(a;q)_\infty (q;q)_\infty}{2\pi i (c;q)_\infty} \lim_{N\to\infty} \oint_{C_N^0} \frac{z^{-\log x/\log q}(cz^{-1};q)_\infty}{(az^{-1};q)_\infty (z;q)_\infty}\, dz. \tag{5.12}$$

In order to estimate the contribution from

$$C_N^1 = \left\{ z = q^{-N-1/2} e^{it} : |t| < \frac{\pi}{2} \right\}, \tag{5.13}$$

we need a bound on the integrand for large $|z|$

$$\sup_{|z|=q^{-N-1/2}} \left| \frac{z^{-\log x/\log q}}{(z;q)_\infty} \right| \leq \left| \frac{|x|^{N+1/2} e^{-\pi^2/\log q}}{(q^{-1/2};q)_\infty \prod_{n=1}^{N}(1-q^{-1/2-n})} \right| = O(q^{N^2/2}), \tag{5.14}$$

as it is dominated by the product in the denominator. From

$$\left| \int_{C_N^1} \frac{z^{-\log x/\log q}(cz^{-1};q)_\infty}{(az^{-1};q)_\infty (z;q)_\infty} dz \right| \leq \pi q^{-N-1/2} \sup_{|z|=q^{-N-1/2}} \left| \frac{z^{-\log x/\log q}(cz^{-1};q)_\infty}{(az^{-1};q)_\infty (z;q)_\infty} \right| = O(q^{N^2/2}), \tag{5.15}$$

and (5.12), we have completed the proof. □



Figure 1. Path of integration C



Figure 2. Path of integration C_N^0

By using Lemma 5.1, it holds that

$$A_1 = \frac{(p;p)_\infty}{2\pi i(-p;p)_\infty} \int_{C_1} \frac{(pa_2tz^{-1};p)_\infty e^{\pi i \log z/\log p}}{z(pa_2{}^{-1}tz^{-1};p)_\infty (z;p)_\infty} dz, \tag{5.16}$$

where the contour C_1 runs from $-i\infty$ to $i\infty$ so that the poles of $1/(z;q)_\infty$ lie to the right of the contour and the other poles lie to the left of the contour. We divide the integral path C_1 into L_1 which runs from $-i\infty$ to $i\infty$ so that all poles lie to the right of the contour and $\hat{C}_1$ which is a anticlockwise-oriented contour and encircles all poles except for the poles of $1/(z;q)_\infty$ (as shown in Figure 3).

(1) Path L_1. Set

$$I_1(s) = \int_{L_1} \frac{(pa_2tz^{-1};p)_\infty e^{\pi i \log z/\log p}}{z(pa_2{}^{-1}tz^{-1};p)_\infty (z;p)_\infty} dz. \tag{5.17}$$



Figure 3. The contours of integration L_1 and $\hat{C}_1$

Using Proposition 4.7, we rewrite $I_1(s)$ as

$$I_1(s) = \int_{L_1} e^{-g_1(z)/\epsilon^2} f_1(z)[1 + O(\epsilon)]\, dz, \tag{5.18}$$

where

$$g_1(z) = Li_2\left(-\frac{2}{z}\right) - Li_2\left(\frac{2}{z}\right) - Li_2(z) + \pi i \log z + is\epsilon \log\left(\frac{z-2}{z+2}\right), \tag{5.19}$$

$$f_1(z) = e^{-2s^2 z/(z^2-4)}(z-2)^{1/2+\alpha}(z+2)^{-1/2-2\alpha}(1-z)^{-1/2}z^{\alpha-1}. \tag{5.20}$$

By the identity of dilogarithm [12]

$$Li_2(z^{-1}) = -Li_2(z) - \frac{(\log z)^2}{2} - \pi i \log z + \frac{\pi^2}{3}, \tag{5.21}$$

$g_1(z)$ is rewritten as

$$g_1(z) = Li_2\left(\frac{z}{2}\right) - Li_2\left(-\frac{z}{2}\right) - Li_2(z) + \pi i \log 2 + \frac{3\pi^2}{2} + is\epsilon \log\left(\frac{z-2}{z+2}\right) \tag{5.22}$$

$$= \pi i \log 2 + \frac{3\pi^2}{2} + \pi s\epsilon - is\epsilon z - \frac{z^2}{4} + O(z^3) \quad (|z| < 1). \tag{5.23}$$

Let us first find a zero point of $g_1'(z)$ (saddle point). Differentiating (5.22) by z and using (4.29), we obtain

$$g_1'(z) = \frac{1}{z}\log\left(1+\frac{z^2}{z-2}\right) + \frac{4is\epsilon}{z^2-4} \tag{5.24}$$

$$= -is\epsilon - \frac{z}{2} - \frac{1+is\epsilon}{4}z^2 + O(z^3) \quad (|z|<1). \tag{5.25}$$

We note here that since

$$\lim_{\epsilon\to 0} g_1'(z) = \frac{1}{z}\log\left(1+\frac{z^2}{z-2}\right), \tag{5.26}$$

all saddle points coalesce at $z=0$ as $\epsilon \to 0$.

Lemma 5.2 *There only exists a saddle point in the domain* $|z| \le |\epsilon|^a$ $(0 < a < 1/3)$ *and that is order one.*

Proof. Since all saddle points are near the origin for sufficiently small ϵ, there exists $a > 0$ such that all saddle points are in the domain $|z| \le |\epsilon|^a$. We note here that we can add the condition $a < 1/3$ without loss of generality. Set

$$A(z) = -g_1'(z) - is\epsilon - \frac{z}{2}. \tag{5.27}$$

Functions $g_1'(z)$ and $A(z)$ are regular in $|z| \le |\epsilon|^a$ and it holds that

$$|g_1'(z)| = \frac{|\epsilon|^a}{2}|1+O(\epsilon^a)| > \frac{|\epsilon|^{2a}}{4}|1+O(\epsilon^a)| = |A(z)|, \tag{5.28}$$

on the circle $|z| = |\epsilon|^a$. Then the statement follows from Rouché's theorem. □

By substituting

$$z = \sum_{k=1}^{\infty} C_k\epsilon^k, \tag{5.29}$$

in (5.25), the saddle point is given by

$$z_1 = -2is\epsilon + 2s^2\epsilon^2 + O(\epsilon^3), \tag{5.30}$$

Since

$$g_1''(z_1) = -\frac{1}{2} + O(\epsilon), \tag{5.31}$$

the steepest descent direction is given as

$$\arg(z-z_1) = \frac{\pi}{2} - \frac{1}{2}\arg\left(\frac{1+O(\epsilon)}{2\epsilon^2}\right) \sim \frac{\pi}{2}. \tag{5.32}$$

In order that L_1 passes in the steepest descent direction, it is necessary that

$$\mathrm{Re}(z_1) = 2\mathrm{Im}(s)\epsilon + 2\left(\mathrm{Re}(s)^2 - \mathrm{Im}(s)^2\right)\epsilon^2 + O(\epsilon^3) < 0. \tag{5.33}$$

Therefore we obtain the following condition

$$\epsilon > 0 \quad (\text{when } \mathrm{Im}(s) < 0), \tag{5.34}$$

$$\epsilon < 0 \quad (\text{when } \mathrm{Im}(s) > 0). \tag{5.35}$$

Now, we consider an approximation of the integrand of $I_1(s)$. We assume that z is in the set

$$D = \left\{ z = \epsilon^k A(\epsilon) \;\middle|\; k \in \mathbb{R}_{>0},\ 0 < |A(0)| < \infty \text{ or } A(\epsilon) = 0 \right\}. \tag{5.36}$$

For simplicity, we first change the variable z to $\hat{z}$ so that the saddle point is given as $\hat{z} = -2is\epsilon$. From (5.31), we set $\hat{z}$ so as to satify

$$g_1'(z)dz = -\frac{1}{2}(\hat{z} + 2is\epsilon)d\hat{z}. \tag{5.37}$$

Moreover, from (5.23) and some technical reasons, we also require

$$g_1(z) = -\frac{\hat{z}^2}{4} - is\epsilon\hat{z} + \pi i \log 2 + \frac{3\pi^2}{2} + \pi s\epsilon + 2is\epsilon^3 + \epsilon^4, \tag{5.38}$$

that is,

$$\hat{z}^2 + 4is\epsilon\hat{z} - 8is\epsilon^3 - 4\epsilon^4 = 4is\epsilon z + z^2 + O(z^3). \tag{5.39}$$

We now add a condition between z and $\hat{z}$ so that the correspondence $z \leftrightarrow \hat{z}$ is $1:1$. To this end, we consider the point $\hat{z} = \hat{w}_0$ corresponding to $z = 0$. From (5.39) $\hat{w}_0$ satisfies

$$\hat{w}_0^2 + 4is\epsilon\hat{w}_0 - 8is\epsilon^3 - 4\epsilon^4 = (\hat{w}_0 - 2\epsilon^2)(\hat{w}_0 + 4is\epsilon + 2\epsilon^2) = 0. \tag{5.40}$$

We here choose $\hat{w}_0 = 2\epsilon^2$, that is, from (5.39) $\hat{z}$ is expressed by z as

$$\hat{z} = -2is\epsilon + 2is\epsilon\sqrt{1 - \frac{2i\epsilon}{s} - \frac{\epsilon^2}{s^2} - \frac{z^2 + 4is\epsilon z + O(z^3)}{4s^2\epsilon^2}}. \tag{5.41}$$

Furthermore, z can be also expressed by $\hat{z}$. Noticing that $z = \epsilon^k A(\epsilon) \in D$, we obtain from (5.39)

$$z = -2is\epsilon \pm 2is\epsilon\sqrt{1 + \frac{2i\epsilon}{s} + \frac{\epsilon^2}{s^2} - \frac{\hat{z}^2 + 4is\epsilon\hat{z} + O\left(A(\epsilon)^3\epsilon^{3k}\right)}{4s^2\epsilon^2}}. \tag{5.42}$$

Since $z = 0$ corresponds to $\hat{z} = \hat{w}_0$, it holds that

$$z = -2is\epsilon + 2is\epsilon\sqrt{1 + \frac{2i\epsilon}{s} + \frac{\epsilon^2}{s^2} - \frac{\hat{z}^2 + 4is\epsilon\hat{z} + O\left(A(\epsilon)^3\epsilon^{3k}\right)}{4s^2\epsilon^2}}. \tag{5.43}$$

For later convenience, we consider the point $z = w_0$ corresponding to $\hat{z} = 0$. By the assumption, w_0 is given in the form

$$w_0 = \epsilon^{k_0} A_0(\epsilon), \tag{5.44}$$

where $k_0 \in \mathbb{R}_{>0}$ and $0 < |A_0(0)| < \infty$. From (5.39), we have

$$\epsilon^{2k_0} {A_0}^2 + 4is\epsilon^{k_0+1} A_0 + 8is\epsilon^3 + 4\epsilon^4 = O(\epsilon^{3k_0}). \tag{5.45}$$

Comparing the degree of ϵ of (5.45), we obtain $k_0 = 1$ or $k_0 = 2$. From (5.43), it holds that

$$w_0 = -2is\epsilon + 2is\epsilon\sqrt{1 + O(\epsilon)} = O(\epsilon^2)\ , \tag{5.46}$$

which indicates

$$k_0 = 2. \tag{5.47}$$

Therefore we can determine w_0 from (5.45) as

$$w_0 = -2\epsilon^2 + O(\epsilon^3). \tag{5.48}$$

Let us next consider the approximation of integrand of $I_1(s)$. We expand $f_1(z)$ in the form

$$f_1(z)\frac{dz}{d\hat{z}} = \sum_{k=0}^{\infty} C_k(\hat{z} - \hat{w}_0)^{k+\rho}. \tag{5.49}$$

Since the characteristic exponent of $f_1(z)$ at $z = 0$ ($\hat{z} = \hat{w}_0$) is $\alpha - 1$ (see (5.20)), it holds that

$$\rho = \alpha - 1. \tag{5.50}$$

Replacing $\hat{z}$ with u by $\hat{z} = 2^{1/2}i\epsilon u$, we obtain

$$I_1(s) = e^{(\pi i \log 2 + \pi^2/2)/\epsilon^2 - \pi s/\epsilon} \sum_{k=0}^{\infty} C_k \int_{L_1} e^{(\hat{z}^2 + 4is\epsilon\hat{z})/4\epsilon^2}(\hat{z} - 2\epsilon^2)^{k+\rho}[1 + O(\epsilon)]d\hat{z} \tag{5.51}$$

$$\begin{aligned} &= (-1)^{(\rho+1)/2} 2^{(\rho+1)/2} \epsilon^{\rho+1} e^{(\pi i \log 2 + \pi^2/2)/\epsilon^2 - \pi s/\epsilon} \\ &\quad \times \sum_{k=0}^{\infty} C_k (-1)^{k/2} 2^{k/2} \epsilon^k \int_{\hat{L}_1} e^{-u^2/2 - 2^{1/2} su}(u + 2^{1/2} i\epsilon)^{k+\rho}[1 + O(\epsilon)]du \end{aligned} \tag{5.52}$$

$$= (-1)^{(\rho+1)/2} 2^{(\rho+1)/2} \epsilon^{\rho+1} e^{(\pi i \log 2 + \pi^2/2)/\epsilon^2 - \pi s/\epsilon} C_0 \int_{\hat{L}_1} e^{-u^2/2 - 2^{1/2} su} u^{\rho}[1 + O(\epsilon)]du, \tag{5.53}$$

where the path of integration $\hat{L}_1$ runs from $-\infty$ to ∞ when $\mathrm{Im}(s) < 0$ and ∞ to $-\infty$ when $\mathrm{Im}(s) > 0$ so that $u = 0$ lies to the right of the path. In the derivetion of (5.53), we note that since $f_1(z)$ does not diverge at $\epsilon \to 0$, C_k also does not diverge. Finally C_0 in (5.53) can be determined in the following manner. From (5.37) and (5.49), we heve

$$-\frac{f_1(z)(\hat{z} + 2is\epsilon)}{2g_1'(z)} = \sum_{k=0}^{\infty} C_k(\hat{z} - 2\epsilon^2)^{k+\alpha-1}. \tag{5.54}$$

Substituting $z = w_0$ ($\hat{z} = 0$) in (5.54), we get

$$C_0 = (-1)^{\alpha+1/2} 2^{-\alpha}. \tag{5.55}$$

Therefore we obtain

$$I_1(s) = (-1)^{3\alpha/2+1/2} 2^{-\alpha/2} \epsilon^{\alpha} e^{(\pi i \log 2 + \pi^2/2)/\epsilon^2 - \pi s/\epsilon} \int_{\hat{L}_1} e^{-u^2/2 - 2^{1/2} su} u^{\alpha-1}[1 + O(\epsilon)]du. \tag{5.56}$$

(2) Path $\hat{C}_1$. Set

$$I_2(s) = \int_{\hat{C}_1} \frac{(pa_2tz^{-1}; p)_\infty e^{\pi i \log z/\log p}}{z(pa_2{}^{-1}tz^{-1}; p)_\infty (z; p)_\infty} dz. \tag{5.57}$$

To evaluate $I_2(s)$, we prepare the following lemma:

Lemma 5.3 *It holds that*

$$\int_C \frac{(cz^{-1};q)_\infty e^{\pi i \log z/\log q}}{z(az^{-1};q)_\infty (z;q)_\infty}\,dz = \frac{2\pi i (ac^{-1};q)_\infty e^{\pi i \log a/\log q}}{(q;q)_\infty (a;q)_\infty}\,{}_2\varphi_1\left(\begin{matrix} a, qac^{-1} \\ 0 \end{matrix}; q, -\frac{c}{a}\right), \tag{5.58}$$

where C is a anticlockwise-oriented contour and encircles all poles except for the poles of $1/(z;q)_\infty$.

Proof. We consider the integral

$$\oint_{C_N} \frac{(cz^{-1};q)_\infty e^{\pi i \log z/\log q}}{z(az^{-1};q)_\infty (z;q)_\infty}\,dz, \tag{5.59}$$

where C_N is a anticlockwise-oriented contour and encircles the poles of $1/(az^{-1};q)_N$. By the residue theorem, we obtain

$$\frac{1}{2\pi i}\oint_{C_N} \frac{(cz^{-1};q)_\infty e^{\pi i \log z/\log q}}{z(az^{-1};q)_\infty (z;q)_\infty}\,dz = \frac{(ac^{-1};q)_\infty e^{\pi i \log a/\log q}}{(q;q)_\infty (a;q)_\infty}\sum_{k=0}^{N-1}\frac{(a;q)_k (qac^{-1};q)_k}{(q;q)_k}\left(-\frac{c}{a}\right)^k. \tag{5.60}$$

Then the statement follows from the limit of $N \to \infty$. ◻

By using Lemma 5.3, $I_2(s)$ is rewritten as

$$I_2(s) = \frac{2\pi i (a_2{}^{-2};p)_\infty e^{\pi i \log(pa_2{}^{-1}t)/\log p}}{(p;p)_\infty (pa_2{}^{-1}t;p)_\infty}\,{}_2\varphi_1\left(\begin{matrix} pa_2{}^{-1}t, pa_2{}^{-2} \\ 0 \end{matrix}; p, -a_2{}^2\right), \tag{5.61}$$

Finally by using the Heine's transformation [20]

$${}_2\varphi_1\left(\begin{matrix} a, b \\ 0 \end{matrix}; q, z\right) = \frac{(az, b;q)_\infty}{(z;q)_\infty}\,{}_2\varphi_1\left(\begin{matrix} z, 0 \\ az \end{matrix}; q, b\right), \tag{5.62}$$

we obtain

$$I_2(s) = \frac{2\pi i (a_2{}^{-2}, -p;p)_\infty e^{\pi i \log(pa_2{}^{-1})/\log p}}{(-a_2{}^2, p;p)_\infty (-1)^{\log t/\log p}}\,{}_2\varphi_1\left(\begin{matrix} -a_2{}^2, 0 \\ -p \end{matrix}; p, pa_2{}^{-1}t\right). \tag{5.63}$$

This completes the proof of Lemma 4.5.

5.2 Proof of Lemma 4.6

We consider the continuous limit of

$$A_2 = {}_2\varphi_1\left(\begin{matrix} -a_2{}^2, 0 \\ -p \end{matrix}; p, pa_2{}^{-1}t\right), \tag{5.64}$$

with (5.6). By using the Heine's transformation [20]

$${}_2\varphi_1\left(\begin{matrix} a, 0 \\ c \end{matrix}; q, z\right) = \frac{1}{(z;q)_\infty}\,{}_1\varphi_1\left(\begin{matrix} a^{-1}c \\ c \end{matrix}; q, az\right), \tag{5.65}$$

A_2 is rewritten as

$$A_2 = \frac{1}{(pa_2{}^{-1}t;p)_\infty}\,{}_1\varphi_1\left(\begin{matrix} pa_2{}^{-2} \\ -p \end{matrix}; p, -pa_2 t\right). \tag{5.66}$$

We prepare a suitable integral representation for A_2.

Lemma 5.4 *It holds that*

$$ {}_1\varphi_1\left(\begin{matrix} a \\ c \end{matrix}; q, x\right) = \frac{(a;q)_\infty (q;q)_\infty}{2\pi i (c;q)_\infty} \int_C \frac{z^{-\log x/\log q}(cz^{-1};q)_\infty}{(az^{-1};q)_\infty (z;q)_\infty}\, dz, \tag{5.67} $$

where the path C runs from $-\infty$ to $+\infty$ so that the poles of $1/(z;q)_\infty$ lie to the right of the path and the other poles lie to the left of it.

The proof of Lemma 5.4 is same as the proof of Lemma 5.1. By using Lemma 5.4, it holds that

$$ A_2 = \frac{({pa_2}^{-2};p)_\infty (p;p)_\infty}{2\pi i(-p;p)_\infty ({pa_2}^{-1}t;p)_\infty} \int_{C_2} \frac{z^{-\log(-pa_2t)/\log p}(-pz^{-1};p)_\infty}{({pa_2}^{-2}z^{-1};p)_\infty (z;p)_\infty}\, dz, \tag{5.68} $$

where the path C_2 runs from $-\infty$ to $+\infty$ so that the poles of $1/(z;q)_\infty$ lie to the right of the path and the other poles lie to the left of it. We divide the path C_2 into L_2 which runs from $-\infty$ to $+\infty$ so that all poles lie to the right of the path and $\hat{C}_2$ which is a anticlockwise-oriented contour and encircles all poles except for the poles of $1/(z;q)_\infty$.

We first consider the path L_2. Set

$$ I_3(s) = \int_{L_2} \frac{e^{-\log(-pa_2t)\log z/\log p}(-pz^{-1};p)_\infty}{({pa_2}^{-2}z^{-1};p)_\infty (z;p)_\infty}\, dz. \tag{5.69} $$

In a similar manner to $I_1(s)$, it holds that

$$ I_3(s) = (-1)^{-3\alpha+1} 2^{-\alpha/2} \epsilon^{-\alpha+1} e^{(-\pi^2/12+\pi i \log 2)/\epsilon^2 - \pi s/\epsilon} \int_{\hat{L}_2} e^{-u^2/2 - 2^{1/2} isu} u^{-\alpha} [1 + O(\epsilon)]\, du, \tag{5.70} $$

under the assumption

$$ \epsilon > 0 \quad (\text{when } \mathrm{Re}(s) < 0), \tag{5.71} $$

$$ \epsilon < 0 \quad (\text{when } \mathrm{Re}(s) > 0). \tag{5.72} $$

Here the path $\hat{L}_2$ runs from $-\infty$ to $+\infty$ when $\mathrm{Re}(s) > 0$ and $+\infty$ to $-\infty$ when $\mathrm{Re}(s) < 0$ so that $u = 0$ lie to the right of the path.

We next consider the path $\hat{C}_2$. Set

$$ I_4(s) = \int_{\hat{C}_2} \frac{e^{-\log(-pa_2t)\log z/\log p}(-pz^{-1};p)_\infty}{({pa_2}^{-2}z^{-1};p)_\infty (z;p)_\infty}\, dz. \tag{5.73} $$

Lemma 5.5 *It holds that*

$$ \begin{aligned} &\int_C \frac{e^{-\log(-pa_2t)\log z/\log p}(-pz^{-1};p)_\infty}{({pa_2}^{-2}z^{-1};p)_\infty (z;p)_\infty}\, dz \\ &\quad = -\frac{2\pi i(-{a_2}^2;p)_\infty e^{2\log(-pa_2t)\log a_2/\log p}}{{a_2}^3 t(p;p)_\infty ({pa_2}^{-2};p)_\infty}\, {}_2\varphi_1\left(\begin{matrix} {pa_2}^{-2}, -{pa_2}^{-2} \\ 0 \end{matrix}; p, \frac{a_2}{t}\right), \end{aligned} \tag{5.74} $$

where C is a anticlockwise-oriented contour and encircles all poles except for the poles of $1/(z;p)_\infty$.

In a similar manner to the proof of Lemma 5.3, we can prove Lemma 5.5. By Lemma 5.5, $I_4(s)$ is rewritten as

$$I_4(s) = -\frac{2\pi i(-{a_2}^2; p)_\infty e^{2\log(-pa_2 t)\log a_2/\log p}}{{a_2}^3 t(p; p)_\infty (p{a_2}^{-2}; p)_\infty}\, {}_2\varphi_1\left(\begin{matrix} p{a_2}^{-2}, -p{a_2}^{-2} \\ 0 \end{matrix}; p, \frac{a_2}{t}\right). \tag{5.75}$$

By using the Heine's transformation [20]

$${}_2\varphi_1\left(\begin{matrix} a, b \\ 0 \end{matrix}; q, z\right) = \frac{(bz; q)_\infty}{(z; q)_\infty}\, {}_1\varphi_1\left(\begin{matrix} b \\ bz \end{matrix}; q, az\right), \tag{5.76}$$

and Lemma 5.4, we obtain

$$I_4(s) = -\frac{(-{a_2}^2, -p{a_2}^{-2}; p)_\infty e^{2\log(-pa_2 t)\log a_2/\log p}}{{a_2}^3 t(p{a_2}^{-2}, a_2 t^{-1}; p)_\infty} \int_{C_3} \frac{e^{-\log(p{a_2}^{-1}t^{-1})\log z/\log p}(-p{a_2}^{-1}t^{-1}z^{-1}; p)_\infty}{(-p{a_2}^{-2}z^{-1}; p)_\infty (z; p)_\infty}\, dz, \tag{5.77}$$

where the path C_3 runs from $-\infty$ to $+\infty$ so that the poles of $1/(z; q)_\infty$ lie to the right of the path and the other poles lie to the left of it. We divide the path C_3 into L_3 which runs from $-\infty$ to $+\infty$ so that all poles lie to the right of the path and $\hat{C}_3$ which is a anticlockwise-oriented contour and encircles all poles except for the poles of $1/(z; q)_\infty$.

We finally consider the paths L_3 and $\hat{C}_3$. Set

$$I_5(s) = \int_{L_3} \frac{e^{-\log(p{a_2}^{-1}t^{-1})\log z/\log p}(-p{a_2}^{-1}t^{-1}z^{-1}; p)_\infty}{(-p{a_2}^{-2}z^{-1}; p)_\infty (z; p)_\infty}\, dz, \tag{5.78}$$

$$I_6(s) = \int_{\hat{C}_3} \frac{e^{-\log(p{a_2}^{-1}t^{-1})\log z/\log p}(-p{a_2}^{-1}t^{-1}z^{-1}; p)_\infty}{(-p{a_2}^{-2}z^{-1}; p)_\infty (z; p)_\infty}\, dz. \tag{5.79}$$

In a similar manner to $I_1(s)$, it holds that

$$I_5(s) = (-1)^{1/2+\alpha} 2^{-3\alpha/2} \epsilon^{1-\alpha} e^{(\pi^2+2(\log 2)^2)/4\epsilon^2 + is\log 2/\epsilon + s^2/2} \int_{\hat{L}_2} e^{-u^2/2 - 2^{1/2} isu} u^{-\alpha} [1 + O(\epsilon)]\, du. \tag{5.80}$$

To evaluate $I_6(s)$, we prepare the following lemma:

Lemma 5.6 *It holds that*

$$\begin{aligned} &\int_C \frac{e^{-\log(p{a_2}^{-1}t^{-1})\log z/\log p}(-p{a_2}^{-1}t^{-1}z^{-1}; p)_\infty}{(-p{a_2}^{-2}z^{-1}; p)_\infty (z; p)_\infty}\, dz \\ &\quad = -\frac{2\pi i t(a_2 t^{-1}; p)_\infty e^{-\log(p{a_2}^{-1}t^{-1})\log(-{a_2}^{-2})/\log p}}{a_2(-p{a_2}^{-2}, p; p)_\infty}\, {}_2\varphi_1\left(\begin{matrix} p{a_2}^{-1}t, -p{a_2}^{-2} \\ 0 \end{matrix}; p, {a_2}^2\right), \end{aligned} \tag{5.81}$$

where C is a anticlockwise-oriented contour and encircles all poles except for the poles of $1/(z; p)_\infty$.

In a similar manner to the proof of Lemma 5.3, we can prove Lemma 5.6. By Lemma 5.6, $I_6(s)$ is rewritten as

$$I_6(s) = -\frac{2\pi i t(a_2 t^{-1}; p)_\infty e^{-\log(p{a_2}^{-1}t^{-1})\log(-{a_2}^{-2})/\log p}}{a_2(-p{a_2}^{-2}, p; p)_\infty}\, {}_2\varphi_1\left(\begin{matrix} p{a_2}^{-1}t, -p{a_2}^{-2} \\ 0 \end{matrix}; p, {a_2}^2\right). \tag{5.82}$$

By using the Heine's transformation (5.62), we obtain

$$I_6(s) = -\frac{2\pi i t(a_2 t^{-1}, -p, p{a_2}^{-1}t; p)_\infty e^{-\log(p{a_2}^{-1}t^{-1})\log(-{a_2}^{-2})/\log p}}{a_2(-p{a_2}^{-2}, p, {a_2}^2; p)_\infty}\, {}_2\varphi_1\left(\begin{matrix} {a_2}^2, 0 \\ -p \end{matrix}; p, p{a_2}^{-1}t\right). \tag{5.83}$$

This completes the proof of Lemma 4.6.

6 Concluding remarks

In this paper, we have constructed the hypergeometric solutions to the symmetric q-Painlevé equations of the types $(A_2+A_1)^{(1)}$, $A_4^{(1)}$, $D_5^{(1)}$, $E_6^{(1)}$, $E_7^{(1)}$ and $E_8^{(1)}$, and discussed their continuous limits. In particular, we have shown that the hypergeometric function appearing in the solution to the symmetric q-P($A_4^{(1)}$) actually reduces to the Weber function by applying the saddle point method to its integral representation.

Before closing, we give a remark on a q-Painlevé equation of the type $A_4^{(1)}$. The following q-difference equation

$$\overline{g}g = \frac{(f+t^{-1})(f+\alpha t^{-1})}{1+\gamma f}, \qquad f\underline{f} = \frac{(g+q^{1/2}\alpha\beta t^{-1})(g+q^{1/2}\beta^{-1}t^{-1})}{1+\gamma^{-1}g}, \tag{6.1}$$

is usually referred to as a q-Painlevé equation of type $A_4^{(1)}$. Equation (6.1) also describes a translation on the root lattice of type $A_4^{(1)}$, but its direction is different from that of (2.5). It may be interesting note that the symmetric q-Painlevé equation obtained from (6.1) by the projective reduction

$$\widetilde{X}\underset{\sim}{X} = \frac{(X+t^{-1})(X+\alpha t^{-1})}{1+X}, \tag{6.2}$$

has no hypergeometric solution which is a sharp contrast to (2.6). Properties of (6.2) will be discussed in detail in a forthcoming paper [26].

Acknowledgement. The authors would like to express their sincere thanks to Mr. P. Howes, Prof. H. Ochiai and Prof. T. Tsuda for fruitful discussions and valuable suggestions. This work has been partially supported by JSPS Grant-in-Aid for Scientific Research No. 23340037 and 22·4366.

References

[1] E. Brézin and V.A. Kazakov, Exactly solvable theories of closed strings, Phys. Lett. B **236** (1990) 144–150.

[2] M.R. Douglas and S.H. Shenker, Strings in less than one dimension, Nucl. Phys. B **335** (1990) 635–654.

[3] A.S. Fokas, A.R. Its and A.V. Kitaev, The isomonodromy approach to matrix models in 2D quantum gravity, Comm. Math. Phys. **147** (1992) 395–430.

[4] B. Grammaticos, F.W. Nijhoff, V. Papageorgiou, A. Ramani and J. Satsuma, Linearization and solutions of the discrete Painlevé III equation, Phys. Lett. A **185** (1994) 446–452.

[5] G. Gasper and M. Rahman, Basic Hypergeometric Series (Second edition), Encyclopedia of Mathematics and Its Applications **96** (Cambridge University Press, Cambridge, 2004).

[6] B. Grammaticos and A. Ramani, On a novel q-discrete analogue of the Painlevé VI equation, Phys. Lett. A **257** (1999) 288–292.

[7] B. Grammaticos and A. Ramani, Discrete Painlevé equations: a review, Lecture Notes in Physics **644** (2004) 245–321.

[8] B. Grammaticos, A. Ramani and V. Papageorgiou, Do integrable mappings have the Painlevé property?, Phys. Rev. Lett. **67** (1991) 1825–1828.

[9] T. Hamamoto, K. Kajiwara and N.S. Witte, Hypergeometric solutions to the q-Painlevé equation of type $(A_1 + A_1')^{(1)}$, Int. Math. Res. Not. **2006** (2006) Article ID 84619.

[10] M.E.H. Ismail and M. Rahman, The associated Askey-Wilson polynomials, Trans. Amer. Math. Soc. **328** (1991) 201–237.

[11] M. Jimbo and H. Sakai, A q-analog of the sixth Painlevé equation, Lett. Math. Phys. **38** (1996) 145–154.

[12] A.N. Kirillov, Dilogarithm identities, Progr. Theoret. Phys. Suppl. **118** (1995) 61–142.

[13] K. Kajiwara, The discrete Painlevé II equation and the classical special functions, in *Symmetries and integrability of difference equations*, eds. by P. Clarkson and F.W. Nijhoff, London Math. Soc. Lecture Note Ser. 255(Cambridge University Press, Cambridge, 1999) 217–227.

[14] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, Hypergeometric solutions to the q-Painlevé equations, Int. Math. Res. Not. **2004** (2004) 2497–2521.

[15] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, Construction of hypergeometric solutions to the q-Painlevé equations, Int. Math. Res. Not. **2005** (2005) 1441–1463.

[16] K. Kajiwara, N. Nakazono and T. Tsuda, Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$, Int. Math. Res. Not. **2011** (2011) 930–966.

[17] K. Kajiwara, Y. Ohta and J. Satsuma, Casorati determinant solutions for the discrete Painlevé III equation, J. Math. Phys. **36** (1995) 4162–4174.

[18] K. Kajiwara, Y. Ohta, J. Satsuma, B. Grammaticos and A. Ramani, Casorati determinant solutions for the discrete Painlevé-II equation, J. Phys. A: Math. Gen. **27** (1994) 915–922.

[19] K. Kajiwara and J. Satsuma, q-difference version of the two-dimensional Toda lattice equation, J. Phys. Soc. Japan 60 (1991) 3986–3989.

[20] R. Koekoek, P.A. Lesky and R.F. Swarttouw, Hypergeometric orthogonal polynomials and their qq-analogues, Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2010.

[21] M.D. Kruskal, K.M. Tamizhmani, B. Grammaticos and A. Ramani, Asymmetric discrete Painlevé equations, Regul. Chaotic Dyn. **5** (2000) 273–280.

[22] T. Masuda, Classical transcendental solutions of the Painlevé equations and their degeneration, Tohoku Math. J. **56** (2004) 467–490.

[23] R.J. Mcintosh, Some asymptotic formulae for q-shifted factorials, Ramanujan J. **3** (1999) 205–214.

[24] M. Murata, H. Sakai and J. Yoneda, Riccati solutions of discrete Painlevé equations with Weyl group symmetry of type $E_8^{(1)}$, J. Math. Phys. **44** (2003) 1396–1414.

[25] N. Nakazono, Hypergeometric τ Functions of the q-Painlevé Systems of Type $(A_2 + A_1)^{(1)}$, *SIGMA Symmetry Integrability Geom. Methods Appl.* **6** (2010).

[26] N. Nakazono and S. Nishioka, Solutions to a q-analog of the Painlevé III equation of type $D_7^{(1)}$, Funkcial. Ekvac. (in press).

[27] S. Nakao, K. Kajiwara and D. Takahashi, On the Multiplicative $\mathrm{dP_{II}}$ and its Ultradiscretization, Kyushu University Ouyorikigakukenkyushukai Houkoku, 9ME-S2, Evolutionary Advancement in Theories of Soliton (1998) 125–130 (In Japanese).

[28] M. Noumi, Painlevé equations through symmetry (American Mathematical Society, Providence, 2004).

[29] Y. Ohta, A. Ramani and B. Grammaticos, An affine Weyl group approach to the eight-parameter discrete Painlevé equation. Symmetries and integrability of difference equations, J. Phys. A: Math. Gen. **34** (2001) 10523–10532.

[30] Y. Ohta, Self-dual structure of the discrete Painlevé equations, RIMS Kokyuroku **1098** (1999) 130–137 (in Japanese).

[31] K. Okamoto, Studies on the Painlevé equations. III. Second and Fourth Painlevé equation, P_{II} and P_{IV}, Math. Ann. **275** (1986) 221–255.

[32] K. Okamoto, Studies on the Painlevé equations. II. Fifth Painlevé equation P_{V}, Japan. J. Math. **13** (1987) 47–76.

[33] V. Periwal and D. Shevitz, Unitary-matrix models as exactly solvable string theories, Phys. Rev. Lett. **64** (1990) 1326–1329.

[34] T. Prellberg, Uniform q-series asymptotics for staircase polygons, J. Phys. A **28** (1995) 1289–1304.

[35] G.R.W. Quispel, J.A.G Roberts and C.J. Thompson, Integrable mappings and soliton equations, Phys. Lett. A **126** (1988) 419–421.

[36] G.R.W. Quispel, J.A.G Roberts and C.J. Thompson, Integrable mappings and soliton equations II, Physica D **34** (1989) 183–192.

[37] A. Ramani and B. Grammaticos, Discrete Painlevé equations: coalescences, limits and degeneracies, Physica A **228** (1996) 150–159.

[38] A. Ramani, B. Grammaticos and J. Hietarinta, Discrete versions of the Painlevé equations, Phys. Rev. Lett. **67** (1991) 1829–1832.

[39] A. Ramani, B. Grammaticos, T. Tamizhmani and K. M. Tamizhmani, Special function solutions of the discrete Painlevé equations. Advances in difference equations, III, Comput. Math. Appl. **42** (2001) 603–614.

[40] A. Ramani,Y. Ohta, J. Satsuma and B. Grammaticos, Self-duality and schlesinger chains for the asymmetric d-P_{II} and q-$\mathrm{P}_{\mathrm{III}}$ equations, Comm. Math. Phys. **192** (1998) 67–76.

[41] H. Sakai, Casorati determinant solutions for the q-difference sixth Painlevé equation, Nonlinearity **11** (1998) 823–833.

[42] H. Sakai, Rational surfaces associated with affine root systems and geometry of the Painlevé equations, Comm. Math. Phys. **220** (2001) 165–229.

[43] K.M. Tamizhmani, B. Grammaticos, A.S. Carstea and A. Ramani, The q-discrete Painlevé IV equations and their properties, Regul. Chaotic Dyn. **9** (2004) 13–20.

[44] R. Wong, Asymptotic approximations of integrals, (Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001).